

OCCASIONAL PAPERS THE MUSEUM TEXAS TECH UNIVERSITY

NUMBER 127

17 MAY 1989

A NEW SUBSPECIES OF THE ROCK SQUIRREL, *SPERMOPHILUS VARIEGATUS*, FROM ISLA TIBURÓN, SONORA, MEXICO

J. KNOX JONES, JR., AND RICHARD W. MANNING

The rock squirrel, *Spermophilus variegatus*, inhabits a broad region in the southwestern United States and adjacent México, north to Colorado, Nevada, and Utah, and southward to Jalisco, Michoacán, and Puebla (Hall, 1981:400). The species has been known for more than a half century from Isla Tiburón, in the Sea of Cortez off central Sonora (Burt, 1938:35), but that population never has been described in detail.

Spermophilus variegatus grammurus (Say, 1823), with type locality on the Purgatory River, Las Animas Co., Colorado, is the subspecies of rock squirrel of mainland Sonora adjacent to Tiburón. The insular population differs from *S. v. grammurus* in several distinctive ways, and represents an hitherto unrecognized race, which is named and described as follows:

Spermophilus variegatus tiburonensis, new subspecies

Holotype.—Adult male, skin and skull, Museum of Vertebrate Zoology no. 97612, from 3 mi. NE Ensenada del Perro, S end Isla Tiburón, Sonora; obtained on 10 January 1942 by M. Delgadillo, original no. 5125 of Seth B. Benson.

Distribution.—Known only from Isla Tiburón, Sonora.

Diagnosis.—Size small to medium among subspecies of species, somewhat smaller than specimens from adjacent mainland of Sonora and elsewhere in range of *S. v. grammurus*. Head, shoulders, and frequently upper back brownish interspersed with few buff-tipped hairs to dark chocolate brown overall, quite in

contrast to relatives on the adjacent mainland (see descriptions of color in Howell, 1938:136-147).

Comparisons.—The new subspecies needs comparison only with *S. v. grammurus*, which is broadly distributed in the American Southwest and adjacent northwestern México, including western Sonora (Hall, *loc. cit.*). From that subspecies, *tiburónensis* differs in being smaller in most cranial dimensions in specimens we have examined (see also measurements of *grammurus* in Hoffmeister, 1986:178-179) and in having a narrower mesopterygoid fossa, smaller auditory bullae, less ovoid (rounder) foramen magnum, less flattened cranium, and a shorter maxillary toothrow. The dark brownish hood and cape immediately separate the new race from *grammurus* and also from the more southerly and easterly *S. v. rupestris*, in which there is no marked darkening of the head and shoulders, as there is, for example, in the blackish-backed *S. v. buckleyi* of the Edwards Plateau region of Texas. All eight specimens examined from Tiburón have the distinctive dark brownish head and shoulders. Otherwise, aside from a darker gular region and, on the average, slightly darker feet and legs, *tiburónensis* generally resembles *grammurus* in color.

Selected external and cranial measurements (in millimeters) of the holotype, followed by those of an adult female (MVZ 74835) and those of two adult females of *S. v. grammurus* (KU 79114 and 80327 from 8 mi. SE Alamos and 9 mi. NNE Imuris, Sonora, respectively) are as follows: total length, 500, 513, 525, 515; length of tail, 222, 230, 244, 216; length of hind foot, 61, 60, 61, 64; length of ear, 25, 25, 28, 27; greatest length of skull, 60.26, 60.05, 63.84, 62.38; condylobasal length, 57.46, 56.60, 60.13, 60.15; rostral breadth, 10.16, 9.67, 10.22, 10.99; postorbital constriction, 16.20, 16.45, 16.13, 16.97; zygomatic breadth, 36.18, 36.86, 39.21,—; mastoid breadth, 21.09, 21.76, 22.55, 23.78; greatest breadth across upper molars, 15.84; 16.51, 17.94, 17.47; breadth of mesopterygoid fossa, 4.03, 4.81, 4.81, 5.41; alveolar length of maxillary toothrow, 12.53, 12.28, 13.26, 12.66; alveolar length of mandibular cheekteeth, 12.32, 11.86, 12.76, 12.48. The adult female from Tiburón weighed 675 grams.

Remarks.—Of the eight specimens available to us from Tiburón, six unfortunately are less than fully adult and thus their measurements cannot be used in direct comparisons. As noted, however, all eight possess the distinctive color pattern of the new subspecies, varying only in degree of brownish

coloration on the head and neck and extent of brownish on the upper back. Tiburón evidently is the only island in the Sea of Cortez on which *Spermophilus variegatus* occurs, one of only 13 native terrestrial, nonvolant mammals known from the island (Lawlor, 1983), none of which is an endemic species. The subspecific name obviously relates to that distribution.

Specimens examined.—Ensenada del Perro, S end Tiburón, 1 (MVZ 97611); 1 mi. W Ensenada del Perro, S end Tiburón, 1 (MVZ 97610); 3 mi. W Ensenada del Perro, S end Tiburón, 1 (MVZ 97609); 3 mi. NE Ensenada del Perro, S end Tiburón, 1 (MVZ 97612); near Las Carrizales, about 2500 ft., about 6 mi. W Bahía Santa Rosa, Tiburón, 1 (MVZ 74835); S end Isla Tiburón, 1 (KU 95282); NE Willard's Pt., Tiburón, 2 (SDSNH 19078-79).

Acknowledgments.—For the loan of specimens, we are grateful to J. L. Patton and B. R. Stein, Museum of Vertebrate Zoology, University of California (MVZ), R. M. Timm, Museum of Natural History, University of Kansas (KU), and A. M. Rea, San Diego Society of Natural History (SDSNH). We also are indebted to Michael A. Bogan and Clyde Jones for critical review of the manuscript.

LITERATURE CITED

- BURT, W. H. 1938. Faunal relationships and geographic distribution of mammals in Sonora, Mexico. Misc. Publ. Mus. Zool., Univ. Michigan, 39:1-77, fold-out map.
- HALL, E. R. 1981. The mammals of North America. John Wiley & Sons, New York, 1:xv + 1-600 + 90.
- HOFFMEISTER, D. F. 1986. Mammals of Arizona. Univ. Arizona Press, Tucson, xx + 602 pp.
- HOWELL, A. H. 1938. Revision of the North American ground squirrels. . . . N. Amer. Fauna, 56:1-256.
- LAWLOR, T. R. 1983. The mammals. Pp. 265-289, 482-494, in Island biogeography in the Sea of Cortez (T. J. Case and M. L. Cody, eds.), Univ. California Press, Berkeley, xi + 508 pp.

Address of the authors: *The Museum and Department of Biological Sciences, Texas Tech University, Lubbock, Texas 79409. Received 5 December 1988, accepted 19 December 1988.*

PUBLICATIONS OF THE MUSEUM
TEXAS TECH UNIVERSITY

Three serials of The Museum of Texas Tech University are published by Texas Tech University Press. Short research studies are published as Occasional Papers, whereas longer contributions appear as Special Publications. Papers of practical application to collection management and museum operations are issued in the Museology series. All are numbered separately and published on an irregular basis.

The preferred abbreviation for citing The Museum's Occasional Papers is *Occas. Papers Mus., Texas Tech Univ.*

Institutional subscriptions (\$19/yr., typically 10 numbers issued per year) are available through Texas Tech University Press, Sales Office, Texas Tech University, Lubbock, Texas 79409-1037. Individuals can purchase separate numbers of the Occasional Papers for \$2.00 each from Texas Tech University Press. Remittance in U.S. currency check, money order, or bank draft must be enclosed with request (add \$1.00 per title or 200 pages of publications requested for foreign postage; residents of the state of Texas must pay sales tax on the total purchase price). Copies of the "Revised checklist of North American mammals north of Mexico, 1986" (Jones *et al.*, 1986, *Occas. Papers Mus., Texas Tech Univ.*, 107:1-22) are available at \$1.25 each in orders of 10 or more.

ISSN 0149-175X



Texas Tech University Press
Lubbock, Texas 79409-1037