Recommended Cattle Analgesic & Anesthetic Agents

Drug **Dosage** Per **Route Duration Notes** 100 lb of Action **Analgesics** 1 gm/ PO or Phenylbutazone 4 mg/kg 24 hours Once daily 500 lb I۷ May be divided twice daily. Flunixin (Banamine) 1.1 - 2.2 mg/kg 0.5 - 1 ml IV slow 24 hours Give for 3 days Actylsalicyclic Acid Give twice daily. 50-100 mg/kg 35 - 70 grains PO 12 hours (Asprin) *Normal 1 bolus is 240 grains* Butorphanol 20 - 30 mg total **Adult Cattle** I۷ This is the TOTAL DOSE for Adult Cattle post-op 2-3 hours (Torbutol) Sedatives/Tranqualizers 0.10 - 0.330.05 - 0.15 ml Cattle can be very sensitive, Xylazine is figured IM mg/kg Xylazine (Rompun) based on 100 mg/ml strength bottle. 0.05 - 0.150.02 - 0.06 ml I۷ Do not mistake it for the 20mg/ml strength. mg/kg **Injectable Anesthetics** 0.10 - 0.330.05 - 0.15 ml Cattle can be very sensitive, Xylazine is figured IM mg/kg Xylazine (Rompun) based on 100 mg/ml strength bottle. 0.05 - 0.15I۷ 0.02 - 0.06 ml Do not mistake it for the 20mg/ml strength. mg/kg Ketamine 0.9 ml I۷ 2 mg/kg

Not Recommended: Pentobarbital (Nembutal), Chlorpromazine (Thorazine), Fentanyl & Droperidol (Innovar-Vet)