



OCCASIONAL PAPERS

Museum of Texas Tech University

Number 341

11 October 2016

MERRIAM'S SHREW (*SOREX MERRIAMI*) IN THE DIET OF A MEXICAN SPOTTED OWL (*STRIX OCCIDENTALIS LUCIDA*) FROM GRANT COUNTY, NEW MEXICO

SIMON P. TYE, KEITH GELUSO, AND MIKE R. FUGAGLI

ABSTRACT

Distributional limits of Merriam's Shrew (*Sorex merriami*) are not well understood across its range in North America due to relatively few records for this widespread species. *Sorex merriami* reaches its southernmost distributional limit in New Mexico, but the species only is known from eight documented localities in the state. We discovered the remains of a single *S. merriami* in a regurgitated pellet of a Mexican Spotted Owl (*Strix occidentalis lucida*) in the Pinos Altos Range, Grant County. This represents the first record of *S. merriami* in the Mogollon Plateau in southwestern New Mexico and the southernmost locality for the species in the United States. This record is best interpreted as a range extension because limited surveys for shrews have been conducted in the region. This record represents the second documentation of Merriam's Shrew as prey for the Mexican Spotted Owl.

Key words: distribution, Merriam's Shrew, Mexican Spotted Owl, Mogollon Plateau, New Mexico, *Sorex merriami*

INTRODUCTION

Distributional limits for Merriam's Shrew (*Sorex merriami*) are not well understood due to relatively few records across its broad geographic range in the United States (Armstrong and Jones 1971). Merriam's Shrew occurs in the western United States from the Sierra Nevada and Cascade Range to western North Dakota, western South Dakota, and western Nebraska (Armstrong and Jones 1971; Hall 1981; Mullican 1984; Merlino et al. 2012). The southernmost distributional limits are from the Mogollon Plateau in Arizona and from the Sacramento Mountains in New Mexico (Hall 1932; Diersing 1979; Hoffmeister 1986). Throughout its range, *S. merriami* has been observed in a variety of

habitats, including dry grasslands, sagebrush steppes, piñon-juniper woodlands, ponderosa pine forests, and mixed coniferous forests (Armstrong and Jones 1971; Findley et al. 1975; Hoffmeister 1986).

In New Mexico, *S. merriami* is known from eight published localities near or in the Manzano, Sacramento, San Juan, Sandia, Sangre de Cristo, Sierra Grande, and Zuni mountains (Findley 1956; Jones 1961; Diersing 1979; George 1990; Hafner and Stahlecker 2002). Frey (2004) suspected the species likely occurs in the Mogollon Plateau in southwestern New Mexico due to the close proximity of a record of

S. merriami in southeastern Arizona. Herein we report the first record of *S. merriami* from the Mogollon Plateau in southwestern New Mexico and discuss the

occurrence of this prey species in the diet of Mexican Spotted Owls (*Strix occidentalis lucida*).

METHODS

We examined 21 regurgitated owl pellets collected from a Mexican Spotted Owl roost located 10 km N, 4 km E Pinos Altos, Grant County, New Mexico. Cranial and mandibular characteristics were used to identify one shrew specimen to species, as we recovered from one of the pellets a single mandible and a nearly complete cranium missing a number of teeth. Another soricid mandible was recovered but there were too many issues in keying it out to be comfortable with identification. To distinguish *S. merriami* from other shrew species, two keys were used (Junge and Hoffmann 1981; Carraway 1995). The single mandible had a well-developed post-mandibular foramen, which narrowed the species to six species of *Sorex* occurring in the United States and Canada (Junge and Hoffmann 1981). Based on the socket size of the 3rd and 4th unicuspid in the partial cranium, the 3rd unicuspid was

larger than the 4th unicuspid, eliminating *S. trowbridgii*, a species that is not known to occur in the region (Junge and Hoffmann 1981). Lastly, the front end of the palate was unusually broad forming a wide triangle (Fig. 22 in Junge and Hoffmann 1981) resulting in identification of the skull as *S. merriami*. For additional support of our identification, we used the key in Carraway (1995) to distinguish between the only two species of *Sorex* in the region with a well-developed post-mandibular foramen, that is, *S. merriami* and *S. arizonae*. Height of coronoid process was > 3.9 mm, measuring 4.12 mm, and ratio of height of the coronoid process to the length of the dentary (6.35 mm) was > 0.54 with a ratio of 0.65, thus again providing further support for identification to *S. merriami*. This specimen was deposited in the Museum of Southwestern Biology (MSB #291774), University of New Mexico, Albuquerque.

RESULTS

The partial cranium and mandible of a *S. merriami* was obtained from a roost of a male Mexican Spotted Owl in a small ravine in the Pinos Altos Range. The ravine had a small perennial stream with grassy banks, and Douglas fir (*Pseudotsuga menziesii*) was the dominant tree in the immediate area. The surrounding forest was predominantly ponderosa pine (*Pinus pon-*

derosa) and Gambel's oak (*Quercus gambelii*). Other remains observed in the pellets included six Mexican Woodrats (*Neotoma mexicana*), six invertebrates, four Botta's Pocket Gophers (*Thomomys bottae*), three voles (*Microtus*), two mice (*Peromyscus*), one Western Harvest Mouse (*Reithrodontomys megalotis*), and one other shrew (unidentifiable mandible, MSB #291775).

DISCUSSION

Owl pellets contain bones and indigestible components of prey, and dissection of pellets is a non-intrusive method that can yield information on regional abundance and distribution of small mammals, including rare and some difficult to sample species (Vernon 1972; Mikkola 1983). Our record of *S. merriami* in the Mogollon Plateau in southwestern New Mexico represents a county record, a new mountain range for this shrew in the state, and southernmost locality for

the species. The closest published locality to our record is from near Rose Peak, Greenlee County, Arizona, which is about 123 km WNW of where our pellet was collected (Hall 1932; Hoffmeister 1986). Rose Peak also is located on the Mogollon Plateau. The previous southernmost record in the United States and closest record in New Mexico was from near Mescalero in the Sacramento Mountains, Otero County, located about 235 km ENE of where the pellet was collected (Diersing

1979). We suggest that our record represents a range extension, and not a range expansion (Frey 2009), as this species likely has always occurred in the vicinity but remained undetected, in part due to difficulty in obtaining records of this uncommon shrew species and a general lack of surveys in the area based on relatively few shrews in regional museums from the region. Examination of additional owl pellets and deployment of pitfall arrays likely will allow researchers to further delineate the distributional limits of Merriam's Shrew, as the limits for this patchily distributed species are still not well understood.

Some species of owls migrate, thus the possibility exists that the *S. merriami* was transported from another area during migration. Spotted Owls, however, generally are considered non-migratory across their range (Gutiérrez et al. 1995). Mexican Spotted Owls are known to shift in elevation $\geq 1,000$ m seasonally and travel relatively short distances of 20 to 50 km in the local region (Gutiérrez et al. 1995). Such limited movement between seasons lends further support that even if the owl captured the *S. merriami* on its wintering range and regurgitated the pellet at its summer breeding locality, the shrew was at most from the Black Range in Sierra County or other parts of the Mogollon Plateau in Catron County, although we find this scenario rather unlikely.

Mexican Spotted Owls inhabit rugged, narrow canyons and high-elevation forests in the southwestern

United States and throughout the Sierra Madre Occidental and Sierra Madre Oriental in Mexico (Gutiérrez et al. 1995; Mullet and Ward 2010). In the southwestern United States, some Mexican Spotted Owls nest in patches of dense trees with high canopy cover in cool areas but forage in more open habitats near roost sites (Willey and van Riper III 2014). Their diet, based on percentage of biomass consumed, primarily consists of woodrats (*Neotoma*), mice (*Peromyscus*), rabbits (*Sylvilagus*), voles (*Microtus*), and Botta's Pocket Gophers, with the remainder including bats, birds, insects, and other small mammals (Ganey 1992; Young et al. 1997; Seamans and Gutiérrez 1999; Block et al. 2005; Ganey et al. 2011; Willey 2013). Although numerous studies have examined the diet of this raptor via pellet analysis, to our knowledge only a single prior study has observed *S. merriami* as a prey item of Spotted Owls (Ganey and Block 2005); thus our observation represents the second record of this prey item for this owl species. Merriam's Shrews also have been observed in the regurgitated pellets of Great Horned Owls (*Bubo virginianus*) and Barn Owls (*Tyto alba*) in California, Nebraska, and Wyoming (Bond 1939; Long and Kerfoot 1963; Merlino et al. 2012). By examining many papers on diets of Spotted Owls throughout their distribution, we conclude that shrews are an uncommon prey item and make up a relatively small percentage of observations and percentage biomass in the diet of this species.

ACKNOWLEDGMENTS

We thank Cody A. Dreier for assistance in the laboratory with dissecting owl pellets and Jon L. Dunnum and Adrienne Raniszewski for museum matters at the Museum of Southwestern Biology, University of New Mexico, Albuquerque. Two anonymous review-

ers improved the content of this paper. Our research was supported, in part, by the Undergraduate Research Fellows program at the University of Nebraska Kearney (UNK), the College of Natural and Social Science at UNK, and Hawks Aloft, Albuquerque, New Mexico.

LITERATURE CITED

- Armstrong, D. M., and J. K. Jones, Jr. 1971. *Sorex merriami*. Mammalian Species 2:1–2.
- Block, W. M., J. L. Ganey, P. E. Scott, and R. King. 2005. Prey ecology of Mexican Spotted Owls in pine-oak forests of northern Arizona. *Journal of Wildlife Management* 69:618–629.
- Bond, R. M. 1939. Observations on raptorial birds in the lava beds-Tule Lake region of northern California. *Condor* 41:54–61.

- Carraway, L. N. 1995. A key to recent Soricidae of the western United States and Canada based primarily on dentaries. Occasional Papers of the Natural History Museum, University of Kansas, Lawrence, Kansas 175:1–49.
- Diersing, V. E. 1979. Noteworthy records of Merriam's Shrew, *Sorex merriami*, from New Mexico. The Southwestern Naturalist 24:708–709.
- Findley, J. S. 1956. Merriam Shrew in New Mexico. Journal of Mammalogy 37:277.
- Findley, J. S., A. H. Harris, D. E. Wilson, and C. Jones. 1975. Mammals of New Mexico. University of New Mexico Press, Albuquerque, New Mexico.
- Frey, J. K. 2004. Taxonomy and distribution of the mammals of New Mexico: An annotated checklist. Occasional Papers, Museum of Texas Tech University 240:1–32.
- Frey, J. K. 2009. Distinguishing range expansions from previously undocumented populations using background data from museum records. Diversity and Distributions 15:183–187.
- Ganey, J. L. 1992. Food habits of Mexican Spotted Owls in Arizona. Wilson Bulletin 104:321–326.
- Ganey, J. L., and W. M. Block. 2005. Dietary overlap between sympatric Mexican Spotted and Great Horned Owls in Arizona. Research Paper RMRS-RP-57WWW. Fort Collins, Colorado. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- Ganey, J. L., J. P. Ward, Jr., and D. W. Willey. 2011. Status and ecology of Mexican Spotted Owls in the Upper Gila Mountains recovery unit, Arizona and New Mexico. General Technical Report RMRS-GTR-256WWW. Fort Collins, Colorado. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station.
- George, S. B. 1990. Unusual records of shrews in New Mexico. The Southwestern Naturalist 35:464–465.
- Gutiérrez, R. J., A. B. Franklin, and W. S. LaHaye. 1995. Spotted Owl (*Strix occidentalis*). The birds of North America, Number 179. The Academy of Natural Sciences, Philadelphia, Pennsylvania, and The American Ornithologists Union, Washington, D.C.
- Hafner, D. J., and D. W. Stahlecker. 2002. Distribution of Merriam's Shrew (*Sorex merriami*) and the Dwarf Shrew (*Sorex nanus*), and new records for New Mexico. The Southwestern Naturalist 47:134–137.
- Hall, E. R. 1932. A new shrew of the *Sorex merriami* group from Arizona. Journal of Mammalogy 13:259–262.
- Hall, E. R. 1981. The mammals of North America. 2nd edition. John Wiley & Sons, New York.
- Hoffmeister, D. F. 1986. Mammals of Arizona. University of Arizona Press and Arizona Game and Fish Department, Tucson, Arizona.
- Jones, C. J. 1961. Additional records of shrews in New Mexico. Journal of Mammalogy 42:399.
- Junge, J. A., and R. S. Hoffmann. 1981. An annotated key to the long-tailed shrews (Genus *Sorex*) of the United States and Canada, with notes on Middle American *Sorex*. Occasional Papers of the Museum of Natural History, University of Kansas, Lawrence 94:1–48.
- Long, C. A., and W. C. Kerfoot. 1963. Mammalian remains from owl-pellets in eastern Wyoming. Journal of Mammalogy 44:129–131.
- Merlino, J. N., A. R. Frohberg, J. Harmon, and K. Geluso. 2012. Least and Merriam's Shrews from Banner County, Nebraska. The Prairie Naturalist 44:60–62.
- Mikkola, H. 1983. Owls of Europe. T. and A. D. Poyser, London, England.
- Mullet, T. C., and J. P. Ward, Jr. 2010. Microhabitat features of Mexican Spotted Owl nest and roost sites in the Guadalupe Mountains. Journal of Raptor Research 44:277–285.
- Mullican, T. R. 1994. First record of Merriam's Shrew from South Dakota. The Prairie Naturalist 26:173.
- Seamans, M. E., and R. J. Gutiérrez. 1999. Diet composition and reproductive success of Mexican Spotted Owls. Journal of Raptor Research 33:143–148.
- Vernon, C. J. 1972. An analysis of owl pellets collected in southern Africa. Ostrich 43:109–123.
- Willey, D. W. 2013. Diet of Mexican Spotted Owls in Utah and Arizona. The Wilson Journal of Ornithology 125:775–781.
- Willey, D. W., and C. van Riper, III. 2014. Home range characteristics of Mexican Spotted Owls in the Rincon Mountains, Arizona. The Wilson Journal of Ornithology 126:53–59.
- Young, K. E., P. J. Zwank, R. Valdez, J. L. Dye, and L. A. Tarango. 1997. Diet of Mexican Spotted Owls in Chihuahua and Aguascalientes, Mexico. Journal of Raptor Research 31:376–380.

Addresses of authors:

SIMON P. TYE

*Department of Biology
University of Nebraska at Kearney
Kearney, NE 68849 USA*

MIKE R. FUGAGLI

*1113 West Street
Silver City, NM 88601*

KEITH GELUSO


*Department of Biology
University of Nebraska at Kearney
Kearney, NE 68849 USA
gelusok1@unk.edu*

PUBLICATIONS OF THE MUSEUM OF TEXAS TECH UNIVERSITY

This publication is available free of charge in PDF format from the website of the Natural Science Research Laboratory, Museum of Texas Tech University (nsrl.ttu.edu). The authors and the Museum of Texas Tech University hereby grant permission to interested parties to download or print this publication for personal or educational (not for profit) use. Re-publication of any part of this paper in other works is not permitted without prior written permission of the Museum of Texas Tech University.

Institutional subscriptions to Occasional Papers are available through the Museum of Texas Tech University, attn: NSRL Publications Secretary, Box 43191, Lubbock, TX 79409-3191. Individuals may also purchase separate numbers of the Occasional Papers directly from the Museum of Texas Tech University.

Series Editor: Robert D. Bradley
Production Editor: Lisa Bradley



ISSN 0149-175X

Museum of Texas Tech University, Lubbock, TX 79409-3191