

# Stem Cell Heterogeneity and Plasticity in Epithelia

Giacomo Donati<sup>1</sup> and Fiona M. Watt<sup>1,\*</sup>

<sup>1</sup>King's College London Centre for Stem Cells and Regenerative Medicine, 28<sup>th</sup> Floor, Tower Wing, Guy's Campus, Great Maze Pond, London SE1 9RT, UK

\*Correspondence: [fiona.watt@kcl.ac.uk](mailto:fiona.watt@kcl.ac.uk)

<http://dx.doi.org/10.1016/j.stem.2015.04.014>

Epithelia cover the surfaces and line the cavities of the body. Recent studies have highlighted the existence of multiple stem cell compartments within individual epithelia that exhibit striking plasticity in response to tissue damage, transplantation, or tumor development. New knowledge about the composition of the epithelial niche and the transcription factor networks that maintain cell identity has provided new insights into the extrinsic and intrinsic regulation of stem cell behavior. In addition new *in vitro* tissue substitutes allow better integration of data from human and mouse models.

## Introduction

Epithelia are one of the four tissue types of the body, the others being connective tissue, muscle, and nerve. Epithelia form the body's interface with the environment and line body cavities, such as the lumen of the intestine. A hallmark of epithelia is that they are constituted of sheets of cells that are tightly packed via specialized cell-cell junctions and adhere to a specialized extracellular matrix, known as the basement membrane (Watt and Fujiwara, 2011). Some epithelia, such as the pulmonary epithelium of the lung, comprise a single cell layer (simple epithelium), while others, such as the lining of the mouth, consist of many cell layers and are known as stratified epithelia. In addition, epithelia may contain adnexal structures, such as glands or hair follicles, which have specific functions.

Epithelia are very heterogeneous (Figure 1). In the skin, the epidermis prevents body dehydration, protects against radiation and penetration by microorganisms, and withstands mechanical stress. Simple epithelia that interface the external environment, such as the lung and the intestinal epithelia, also act as a barrier against pathogens in addition to having absorptive functions. Examples of epithelial cells with specific secretory functions are epidermal sebocytes, which release lipids to lubricate the skin surface, and intestinal Paneth cells, which release antimicrobial compounds that are important in host immunity. While Paneth cells are distributed individually within the crypts of the small intestine, sebocytes are organized in glands associated with the hair follicles. In stratified epithelia the stem cells are attached to the basement membrane, while in simple epithelia the differentiated cells, like the stem cells, are attached to the basement membrane.

The epidermis and intestine are two of the tissues in which the existence of stem cells was first postulated. This is because the terminally differentiated cells are unable to divide and must be replaced throughout adult life by less differentiated cells, the stem cells, that both self-renew and generate differentiated progeny (Hall and Watt, 1989). In other epithelia, such as the lung and mammary gland, the kinetics of cell differentiation and replacement are somewhat different. For example, in mammary gland periods of dormancy are interspersed with phases of hormone-driven morphogenesis, at puberty and during pregnancy (Visvader and Stingl, 2014). The rate of cell turnover is not only highly variable between different epithelial tissues—




ranging from a few days in the small intestine crypts to a few weeks in the epidermis and several months in the urinary tract epithelium—but also varies within a single epithelium, with some cells dividing much more rapidly than others. This complexity highlights the tight regulation of stem cell activity required to maintain tissue homeostasis.

In this Review we discuss recent advances in our understanding of epithelial stem cells and the intrinsic and extrinsic factors that regulate their behavior. Within each epithelium multiple stem cell populations are physically and functionally compartmentalized, yet they can exhibit striking plasticity when perturbed. Clonal analysis has revealed epithelial stem cells and their progeny to exhibit dynamic behavior, which reflects in part responses to extrinsic signals from the stem cell niche and in part intrinsic control of cell state via transcriptional and epigenetic regulators. The integration of single-cell data with *in vivo* lineage tracing is increasing our understanding of epithelial stem cell properties, while the generation of epithelia from pluripotent stem cells is blurring the traditional boundary between pluripotent and tissue stem cells.

## Individual Epithelia Contain Multiple Stem Cell Populations

Over the years a range of *in vitro* and *in vivo* assays have been developed to identify and quantify epithelial stem cells (Kretzschmar and Watt, 2012). Of these, lineage-tracing experiments represent the most powerful method to study stem cells under homeostatic conditions. Not only does lineage tracing avoid the need to disaggregate the tissue to identify stem cells, but it also allows visualization of the contribution of stem cell progeny to multiple differentiated cell types, which are often organized into complex adnexal structures such as the hair follicles of the skin.

One important principle to emerge from lineage tracing in multiple epithelia is that of stem cell compartmentalization. It is evident that the intestine, mammary glands, lungs, and epidermis are not obligatorily maintained by a single, homogeneous stem cell population. Rather, there are several stem cell pools, characterized by differences in the molecular markers they express and their context-dependent functions. For example, in mouse skin (Figures 2A and 2B), stem cell populations have been identified in the interfollicular epidermis, hair follicle bulge, hair follicle junctional zone, and sebaceous gland

Epithelia	Cell types	Structure	Adenexa	Selected stem cell markers
<b>Epidermis &amp; pilosebaceous unit</b>				
	<ul style="list-style-type: none"> <li>○ Differentiated cells</li> <li>● Differentiating cells</li> <li>⊖ IFE stem cells</li> <li>⊖ Quiescent stem cells</li> <li>⊖ Matrix cells (HF lineages)</li> <li>⊖ Lgr5 stem cells</li> </ul>	Stratified epithelium	Hair follicle Sweat gland Sebaceous gland	Krt14, 5 Lrig1 Lgr5 Lgr6 Cd34 / Cd49f
<b>Small intestine (villus &amp; crypt)</b>				
	<ul style="list-style-type: none"> <li>○ Absorptive cells</li> <li>○ Goblet &amp; Enteroendocrine cells</li> <li>○ Dll1 cells</li> <li>○ Lgr5 stem cells</li> <li>○ Paneth cells</li> </ul>	Simple epithelium	No	Lgr5 Lrig1 Bmi1 HopX
<b>Distal lung (bronchioli &amp; alveoli)</b>				
	<ul style="list-style-type: none"> <li>○ Ciliated cells</li> <li>○ Clara cells</li> <li>○ Bronchioalveolar stem cells</li> <li>○ Alveolar type 1 cells</li> <li>○ Alveolar type 2 cells</li> </ul>	Simple epithelium	No	Scgb1a1 Scgb1a1 / Sftpc Krt5, 14, 8 Trp63
<b>Mammary gland</b>				
	<ul style="list-style-type: none"> <li>○ Alveolar epithelial cells</li> <li>○ Ductal epithelial cells</li> <li>○ Myoepithelial cells</li> </ul>	Simple epithelium surrounded by myoepithelial cells	No	Krt14, 5, 8 Elf5 Cd49f / Cd29 / Cd24 (EpCAM, Sca1 negative)

**Figure 1. Different Types of Epithelia**

Schematic summary of the architecture and stem cell compartments of different types of epithelia.

(Watt, 2014). In the lung, lineage tracing of Scgb1a1-positive Clara cells suggests a model in which trachea, bronchioles, and alveoli are renewed by distinct epithelial stem cells during adult homeostasis (Rawlins et al., 2009). In mouse esophageal epithelium there are basal cell subpopulations with distinct differentiation and proliferation properties, consistent with a hierarchical model that involves stem cell progeny differentiating in the basement membrane niche (DeWard et al., 2014).


In the intestinal epithelium *Lgr5*-expressing stem cells give rise to multiple differentiated progeny, including absorptive enterocytes and goblet, Paneth, and enteroendocrine cells (Barker et al., 2007) (Figure 2C). Although several studies have indicated the existence of stem cell heterogeneity (Sangiorgi and Capecchi, 2008; Takeda et al., 2011; Li and Clevers, 2010; Itzkovitz et al., 2012; Muñoz et al., 2012), live imaging has recently established that *Lgr5*-positive cells are at the top of the cellular hierarchy in the intestine. Nevertheless, cells located above the crypt base can occasionally revert to the bottom of the crypt after cell division (Ritsma et al., 2014) (Figure 2D).

While the stem cells of individual epithelia are undoubtedly heterogeneous, some stem cell markers are common to multiple epithelia. This has become evident from the development of

transgenic mice expressing Cre recombinase under the control of specific gene promoters. One of the first examples was *Lgr5*, originally found as a stem cell marker in the small intestine, then subsequently found to mark stem cell populations in the lower hair follicle, mammary gland, and stomach (Barker et al., 2007; Jaks et al., 2008). *Lrig1* is also expressed in stem cells of a variety of epithelia, and there is partial overlap of expression between *Lgr5* and *Lrig1* in colon epithelium (Powell et al., 2012). It is likely that the existence of common stem cell markers in different epithelia reflects the fact that common pathways regulate the stem cell compartments, including *Wnt*, *Egf*, and *Notch* (Estrach et al., 2008; Jensen and Watt, 2006; Lim et al., 2013; Schuijers et al., 2015; van Es et al., 2012). For example, *Lgr5* is a *Wnt* target gene and *Lrig1* is a negative regulator of *Egf* signaling.

#### Quantitative Analysis of Epithelial Clones Reveals Dynamic Stem Cell Behavior in Healthy Tissue and in Cancer

Statistical analysis of the size and distribution of epithelial cell clones marked by lineage tracing has given new insights into the behavior of epithelial stem cells. Clone size is relatively



**Figure 2. Epithelial Stem Cell Contributions during Tissue Homeostasis and Repair**

(A and B) Epidermis. (C and D) Intestinal epithelium. (A and B) Under steady-state conditions (homeostasis) distinct epithelial stem cells give rise to distinct differentiated progeny. However, during repair of tissue damage they acquire plasticity and exhibit wider differentiation potential. (A) Lgr5-positive stem cells and their progeny (red), (B) Lrig1-positive stem cells and their progeny (blue). Lgr5- and Lrig1-positive stem cell populations are located in separate HF compartments (lower bulge and isthmus, respectively) and in homeostasis they maintain the lower follicular bulb and the upper HF (infundibulum), respectively. During repair these restrictions are broken and they contribute to re-epithelialization more widely. (C) Lgr5-positive stem cells (green) at the base of the crypts give rise to all the lineages of the intestinal epithelium (left panel). (D) Dll1-positive cells (orange) are progenitors of the secretory lineages in homeostatic epithelium, but upon crypt damage Dll1 cells can revert to Lgr5 stem cells and repopulate the entire crypt.

easy to quantify in epithelia, because intercellular adhesive junctions maintain stem cell progeny in proximity to one another. In mouse tail epidermis, inducible genetic labeling first allowed long-term analysis of the fate of individual cells. Clone size distributions suggested that only one type of cell maintains epidermal homeostasis through both symmetric and asymmetric division (Clayton et al., 2007). This view was subsequently challenged by clonal analysis of cells expressing K14, an epidermal basal layer marker, and *Involucrin*, which is primarily expressed by differentiating cells. The analysis suggested that slow-cycling K14-expressing stem cells give rise to *Involucrin*-positive


transit-amplifying cell progeny (Mascré et al., 2012). In fact the situation is more complex because of the existence of two distinct differentiation programs within tail interfollicular epidermis, which are maintained by discrete cell populations in the basal layer (Gomez et al., 2013). At birth the tail epidermis comprises a single differentiated lineage, but from P9, a second type, known as the scale lineage, is established. The scale expands by more rapid proliferation of the stem cell compartment than that of the interscale, and the relative size of each compartment is controlled by Wnt and Lrig1 signaling (Gomez et al., 2013).

Long-term fate mapping of clones in intestinal epithelium using the R26R-Confetti allele has revealed that stem cells compete to colonize each crypt through neutral stochastic drift dynamics. Intestinal crypts become monoclonal within 1 to 6 months, indicating that there is a natural predisposition toward fewer but larger clones (Ritsma et al., 2014; Snippert et al., 2010) (Figure 3A). Live imaging has shown that individual Lgr5-expressing stem cells at the base of the crypt have a positional advantage in self-renewal ability compared to stem cells in the upper part of the crypt, since dividing cells can push neighboring stem cells out of the niche (Ritsma et al., 2014) (Figure 2C). Multicolor fate mapping in mammary glands suggests that, over time, a few stem cell clones prevail and colonize large areas of the epithelium in a similar manner to the intestine (Rios et al., 2014).

One situation in which lineage analysis has been performed in human tissue is in the intestinal crypt, where somatic mtDNA mutations allow tracing of clonal lineages. This has shown that the clonal evolution of human intestinal stem cells resembles that of the murine crypt and although human crypts house 10 times more cells than mouse, both human and mouse crypts have a similar number of stem cells (Baker et al., 2014).

Mathematical modeling and statistical analysis of in vivo stem cell behavior has not only provided new insights into epithelial homeostasis but can also aid interpretation of the consequences of perturbing homeostasis (Blanpain and Simons, 2013). For example, inhibition of the Notch pathway in esophageal epithelial stem cell clones generates an imbalance in neutral drift dynamics in favor of positive selection of mutant clones at the expense of wild-type clones. This is the result of an increase both in the proliferation rate of the mutant stem cell clones and in their ability to accelerate neighboring wild-type cell differentiation (Alcolea et al., 2014). A similar effect is observed when oncogenic K-ras is sporadically activated in the intestinal LGR5 stem cells. Cell proliferation increases, creating a biased drift toward crypt clonality (Snippert et al., 2014).

Another example of clonal perturbation occurs in “field cancerization:” in the early stages of tumor development an unequal competition between normal and mutant cells leads to clonal expansion of histologically normal but genetically altered cells. Indeed clonal selection is a key process in tumor growth; cancer evolves through the expansion of the most successful genetically altered clones that are able to adapt and proliferate in tumor niches (Figure 3B). The analysis of changes in the size distribution of p53 mutant clones in chronically UVB-irradiated mouse epidermis has revealed that, as in normal epidermis, the fate of individual clones is stochastic, resulting in a balance between cell loss and proliferation. Preneoplastic clones are not derived from long-lived mutant stem cells but from mutant progenitors



**Figure 3. Clonal Heterogeneity in Homeostasis and Cancer**

(A) Multi-color genetic labeling of epithelial cells in the intestine shows neutral competition between stem cells. Over time the number of clones decreases while the clone size increases.

(B) Concept that the mechanisms of stem cell plasticity during tissue repair contribute to the evolutionary dynamics of cancer stem cell clones. Upper panel: clonal evolution resulting in clonal heterogeneity over time. Lower panel: in tumors clones of cancer stem cells with genomic mutations that confer a growth advantage will expand. In addition, changes in the niche (asterisk) could promote clonal evolution without genetic mutation by conferring cell plasticity, as occurs during repair of healthy tissue.

with a random fate (Klein et al., 2010). In mouse intestine the fate of stem cells that have a potential competitive advantage through *Apc* loss, *K-ras* activation, and *P53* mutation has been quantified. Surprisingly, many of the mutant stem cells are stochastically replaced by wild-type stem cells; only *P53* mutations provide a competitive advantage (Vermeulen et al., 2013). These studies raise the interesting question of whether the drift toward monoclonality that is characteristic of normal epithelia increases the likelihood that mutant clones will expand to become tumors (Figure 3B).

Lineage tracing has provided evidence for the existence of cancer stem cells in epithelial tumors. By definition, cancer stem cells have the highest potential to propagate and maintain tumor growth when compared with other cells in a tumor and they represent the top of a cancer cell hierarchy (Figure 3B). In benign epidermal tumors, named papillomas, multiple genetically labeled tumor cells contribute to the bulk of the tumor; however, in malignant squamous cell carcinomas, clonal analysis indicates that tumor formation reflects expansion of individual cancer stem cells (Driessens et al., 2012). In mouse intestine, *Lgr5* cells that represent 5%–10% of all tumor cells are responsible for the growth of established adenomas (Schepers et al., 2012). Lineage tracing has also shown the existence of a small population of cancer stem cells in mammary tumors, although intravital imaging has recently highlighted the dynamic nature of these cells during tumor growth (Zomer et al., 2013).

### Epithelial Stem Cells Exhibit Plasticity during Tissue Repair and Regeneration


In addition to the concept that individual epithelia harbor multiple stem cell populations (stem cell heterogeneity), a second princi-

ple that has emerged is that individual epithelial stem cell populations exhibit wider differentiation potential in tissue reconstitution assays than in the intact tissue (stem cell plasticity; Figures 2A, 2B, and 4). This has been shown for the epidermis, intestinal epithelium, lung, and mammary gland (Van Keymeulen et al., 2011). For instance *Lrig1*-positive epidermal stem cells give rise to differentiated cells that are restricted to specific areas of the pilosebaceous unit under steady state conditions, but in skin reconstitution assays *Lrig1*-positive cells contribute to all the different epidermal compartments (Jensen et al., 2009; Page et al., 2013) (Figure 4A). Even in the hematopoietic system, where serial reconstitution of the blood by individual cells has long been the gold standard for stem cell identification (Rossi et al., 2008), it is reported that cells behave differently under steady-state conditions. This was demonstrated in native non-transplant hematopoiesis by cell labeling in situ through inducible transposon tagging (Sun et al., 2014) and by inducible genetic labeling of *Tie2*-positive hemopoietic stem cells in bone marrow (Busch et al., 2015).

While cell transplantation assays have revealed a broader plasticity of epithelial stem cells than they exhibit under steady-state conditions (Prater et al., 2014; Van Keymeulen et al., 2011), it could be argued that this is of limited physiological relevance. A single mammary gland cell will never normally be found in isolation in a cleared mammary fat pad, and the skin is never normally reconstituted from a mixture of disaggregated epidermal and dermal cells. Nevertheless, the plasticity revealed by those studies is also seen during tissue repair (Figures 2 and 4A). In the hair follicle, *Lgr5*- and *Lrig1*-expressing epidermal stem cells produce differentiated progeny that are normally restricted to the hair follicle. However, after skin wounding their progeny are able to migrate into the interfollicular epidermis where they persist for long periods (Ito et al., 2005; Jaks et al., 2008; Page et al., 2013) (Figures 2A and 2B). Even targeted laser ablation of individual stem cells can provoke a change in cell fate, as in the hair follicle, where cells that have left the stem cell compartment are able to de-differentiate and become bulge stem cells (Rompolas et al., 2013).

In the small intestine genetic ablation of *Lgr5*-positive stem cells does not perturb epithelial homeostasis because of compensation by a different stem cell pool comprising *Bmi1*-expressing cells (Tian et al., 2011). *Dll1*-expressing cells that are committed to the secretory lineage can revert to an intestinal stem cell state after tissue damage (van Es et al., 2012). Similarly it has been shown that mouse intestinal quiescent cells that are committed precursors of secretory Paneth cells and the enteroendocrine lineage can give rise to all the main epithelial cell types after intestinal injury (Buczacki et al., 2013). In the lung, type II pneumocyte progenitors mediate highly focal lung repair, while *p63/Krt5*-positive distal airway stem cells are essential for lung regeneration after acute lung damage (Hogan et al., 2014; Zuo et al., 2015).

The interaction between neighboring epithelial cells in different differentiation states can regulate stem cell behavior and plasticity. In the epidermis, by promoting *Shh* signaling, cells that have exited the stem cell compartment directly stimulate quiescent stem cells to expand and indirectly, via dermal factors, favor their own proliferation (Hsu et al., 2014). When airway stem cells in the lung are ablated, luminal secretory cells can de-differentiate



**Figure 4. Stem Cell Plasticity in Tissue Reconstitution Assays**

(A) In undamaged epidermis Lgr5 (red) and Lrig1 (blue) stem cells contribute to different regions of the tissue. However, when disaggregated, combined with fibroblasts, and implanted into a wound, both stem cell populations can reconstitute all the lineages of the epidermis. (B) Lgr5-positive intestinal stem cells (green) are sufficient to generate crypt-villus units in culture.

Transdifferentiation also occurs in the context of disease, in particular the phenomenon of metaplasia: the replacement of a normal cell type with another differentiated cell type. This phenomenon, although rare, affects multiple epithelia and can predispose an individual to cancer development (Slack, 2007). For example, Barrett's esophagus is a condition in which the normal esophageal stratified squamous epithelium is replaced by a simple columnar epithelium containing goblet cells that are normally found in the intestinal tract.

into basal stem cells that are multipotent and have the ability to repair the epithelium. This de-differentiation process is normally inhibited by cell-cell interactions with basal stem cells (Tata et al., 2013).

Wound healing repairs tissue without necessarily restoring full functionality. In contrast, regeneration is a process that leads to a full recovery of tissue functions. Although adult mammals cannot regrow entire amputated limbs, the distal tips of the digits in mice can be regenerated. The process involves coordinated changes in a wide range of different cell types and stem cell populations without conversion between different lineages (Rinkevich et al., 2011). In the nail, basal epidermal stem cells orchestrate the regeneration process through *Wnt* activation (Takeo et al., 2013).

One emerging concept is that the nature of an injury can determine whether or not tissue is fully regenerated in addition to being repaired. For instance, the excision of a small area of adult mouse skin is repaired through a wound healing process leading to scar formation without de novo formation of hair follicles. However the excision of a large area of skin leads to de novo hair follicle regeneration as a result of *Wnt* activation (Ito et al., 2007). A broad concept that is starting to emerge is that the changes in cell state that occur in response to injury, and therefore the degree to which tissue function is regenerated, depend on the nature and extent of the injury. This could, for example, resolve the current controversy about the cell types that are responsible for liver repair (Grompe, 2014; Schaub et al., 2014; Tarlow et al., 2014).

Transdifferentiation—conversion from one lineage to another—is a well-documented developmental process. For example, during tooth development a significant proportion of mesenchymal stem cells are derived from peripheral nerve-associated glial cells (Kaukua et al., 2014). In addition, epithelial cells can undergo transdifferentiation during tissue repair. In adult pancreas, loss of beta cells can lead to transdifferentiation of acinar cells into beta cells (Thorel et al., 2010). During liver regeneration, hepatocytes can be converted into biliary epithelial cells (Yanger et al., 2013).


Although this is a reversible process, it is associated with development of esophageal cancer. A further example is in breast cancer, where transdifferentiation of mammary epithelial cells into squamous epithelial carcinoma cells reflects a transition from a simple mammary epithelium to a stratified epithelium (Li et al., 2003).

The existence of cancer stem cells and the evolution of cell clones via genetic mutations represent two well-established and interconnected features of cancer (Figure 3B). We hypothesize that a third mechanism that could contribute to clonal expansion and heterogeneity is via acquisition of the same type of cell plasticity as observed during repair of healthy tissue in the absence of genetic mutational events (Figure 3B). This begs the question: what is the nature of stem cell plasticity? At present we do not know whether, within a given epithelium, different stem cell types share a common state of plasticity, and whether the stem cells of a tissue in which homeostasis has been reestablished are the same as, or different from, the stem cells of the undamaged tissue (Figure 5).

#### Fibroblasts, Immune Cells, and Bacteria Are Dynamic Constituents of the Stem Cell Niche

It is well established that the stem cell microenvironment, or niche, regulates many aspects of cell fate. Several generic constituents of the niche, including extracellular matrix composition, physical forces, and neighboring cells, are well known to affect epithelial stem cell behavior (Lane et al., 2014). In addition, three other aspects of the niche are becoming better understood. These are connective tissue fibroblasts, cells of the immune system, and the microbiome. These niche components help to maintain the functional diversity of epithelial stem cells during homeostasis, during tissue repair, and in cancer (Figure 6).

In skin, different mesenchymal cell types regulate epidermal stem cell properties. These include the adipocyte lineage, which positively regulates hair follicle stem cell activity (Festa et al.,




**Figure 5. Model of Stem Cell Plasticity during Repair of Injured Tissue**

Three different stem cell populations are indicated by red, orange, and green circles. The plastic state induced by injury (rectangle) could either be common to all stem cell types (stripes) or different for different stem cell types. Once homeostasis is reestablished, stem cells could revert to their original state (red, yellow, or green circles) or acquire a new identity (multi-colored circles). Alternatively, only a subset of stem cells (yellow circle) could contribute to the repaired tissue while the others undergo cell death.

2011), and the cluster of cells at the base of the hair follicle, known as the dermal papilla, which contributes to maintaining hair follicle identity. Recent studies have shown that different subsets of skin fibroblasts arise from different lineages during development. The upper dermal lineage not only gives rise to the dermal papilla, but also to the arrector pili muscle and papillary fibroblasts. The lower dermal lineage gives rise to reticular fibroblasts involved in the production of the fibrillar extracellular matrix and the adipocyte lineage. The upper lineage is required for hair follicle formation, while the lower lineage mediates the first wave of dermal repair after wounding (Driskell et al., 2013). Epidermal *Wnt* signaling regulates the different dermal compartments (Driskell et al., 2013) by a combination of short (Fujiwara et al., 2011) and long (Donati et al., 2014) range signaling.

The stromal cells of mouse lung also comprise different cell lineages, with different populations being organized into localized and tightly regulated domains within the tissue. As in the epidermis, the different fibroblast subsets regulate epithelial proliferation and reciprocal communication is mediated by *Wnt* signaling (Kumar et al., 2014). Therefore, the picture that emerges both in the epidermis and the lung is that the mesenchyme is just as complex and compartmentalized, in terms of cellular heterogeneity, as the epithelium.

Fibroblast heterogeneity not only contributes to maintaining epithelial heterogeneity during homeostasis but also contributes to cancer progression, since tumor stroma is a key factor in cancer initiation, growth, and progression. Fibroblast secretion of extracellular matrix proteins, such as fibrillar collagens, influences the mechanical properties of connective tissue. In the mammary gland increased matrix stiffness can promote tumor progression through microRNA-dependent PTEN expression (Mouw et al., 2014), suggesting a role for the fibroblast lineages responsible for ECM deposition. The isolation of cancer-associated fibroblasts (CAFs) from different stages of breast cancer



**Figure 6. Fibroblasts, Immune Cells, and Bacteria as Components of the Epithelial Stem Cell Niche during Homeostasis, Tissue Repair, and Cancer**

Two different epithelial compartments within one tissue are shown (pale and dark pink). Each type of stem cell resides in a different niche (bacteria, brown or yellow; fibroblasts, white or gray; and immune cells, green or orange). Double-headed arrows indicate crosstalk between stem cells and niche cells. In homeostasis the different niche constituents may reinforce different stem cell identities. In tissue repair and cancer the different niches may influence stem cell behavior, such as proliferation and cell migration.

progression has shown that CAFs require the transcription co-activator YAP to promote matrix stiffening, cancer cell invasion, and angiogenesis (Calvo et al., 2013). Other signaling pathways and transcription factors that regulate the properties of tumor stroma have also been identified. In skin UVA light downregulates stromal Notch signaling, and mesenchymal-specific deletion of a key Notch effector triggers stromal atrophy, inflammation, and field cancerization (Hu et al., 2012). The transcriptional regulator heat shock factor 1 (HSF1) is frequently activated in CAFs. HSF1-expressing CAFs support malignancy in a non-cell-autonomous way, secreting TGF- $\beta$  and SDF1 (Scherz-Shouval et al., 2014). While the signaling pathways that mediate

the contribution of CAFs to cancer are increasingly well characterized, the possibility that CAFs are heterogeneous rather than a uniform cell population is only just starting to emerge (Öhlund et al., 2014; Sugimoto et al., 2006). The functional heterogeneity of fibroblasts observed in healthy epithelia also contributes to the heterogeneity of CAFs in melanoma (Rinkevich et al., 2015).

Cells of the immune system are key components of epithelia, both in health and in disease (Pasparakis et al., 2014). In the epidermis  $\gamma\delta$  T cells influence survival, proliferation, and migration of keratinocytes and represent a crucial component for hair follicle neogenesis after skin wounding, indicating their role in the epidermal stem cell niche (Gay et al., 2013). As in the case of epithelial-fibroblast communication, communication with immune cells is reciprocal. For example, adult thymic epithelial cells regulate the fate of hematopoietic cells, facilitating the selection of thymocytes with functional T cell receptors (Anderson and Takahama, 2012).

Finally, while it is well known that as part of their barrier function epithelia are colonized by commensal bacteria, it has only recently become apparent that their impact can be on specific epithelial stem cell compartments, particularly in the lung, the skin, and intestine. This is because different microbiota colonize different sites within an epithelium, dependent on environmental factors such as temperature and humidity (Grice and Segre, 2011). The interdependence between epithelial tissue functionality and specific microbiota has been identified. In early life, lung microbiota promote tolerance to allergens. Following exposure to allergens, there is a positive correlation between an increased number of goblet cells and a change of microbiota at the phylum level (Gollwitzer et al., 2014). In the intestinal crypt, Lgr5-positive stem cells express high levels of the cytosolic innate immune sensor Nod2. Nod2 stimulation protects against oxidative stress and promotes cell survival, suggesting that gut repair can be supported by the presence of bacteria (Nigro et al., 2014).

The presence of specific bacteria can also impact epithelial tumor formation. While this is well known in the case of *Helicobacter pylori* and gastric cancer (Sigal et al., 2015), it also appears to be true in the skin. Chronic skin inflammation and wounding can trigger cancer and in a mouse model this has been linked to the presence of flagellated bacteria, which are recognized by Toll-like receptor 5 (Hoste et al., 2015). Pathogen surveillance involves a precise cascade of cellular events in epithelial tissue. For example in skin once Cd103 dendritic cells sense the presence of *Staphylococcus epidermidis*, they migrate to the lymph nodes where they prime Cd8 T cells. Subsequently, commensal specific Cd8 T cells move to the skin where they enhance the anti-microbial defense of keratinocytes (Naik et al., 2015). Intriguingly, dermal adipocytes have recently been shown to participate in skin defense against bacteria, demonstrating that microorganisms are not exclusively part of the epidermal stem cell niche (Zhang et al., 2015).

### Transcription and Epigenetic Factors Regulate Epithelial Stem Cell State

Stem cell identity and function depends on both extrinsic factors (components of the niche) and cell-intrinsic factors. The intrinsic regulators are sequence-specific transcription factors and epigenetic factors controlling transcriptional networks to drive tissue-


specific gene expression patterns. In embryonic stem cells so-called super-enhancers, consisting of large clusters of enhancers bound by chromatin and transcription factors, have been shown to define cell identity (Hnisz et al., 2013). Recently super-enhancers have also been characterized in adult epidermal cells. These TF-dense DNA regions are very dynamic, since they are remodeled in response to microenvironmental changes and differentiation status and during wound healing (Adam et al., 2015).

In epithelia there are several examples of how loss of a single transcription factor can lead to stem cell fate alteration. In the epidermis LHX2 plays a critical role in organizing the niche of hair follicle stem cells, balancing cell quiescence and lineage selection (Folgueras et al., 2013). Loss of the transcription factor Cdx2 leads to transformation of intestinal stem cells into gastric stem cells (Simmini et al., 2014). One explanation for the profound effects of individual transcription factors is that they represent a crucial outcome of cell signaling. In intestinal epithelium, the mechanism that links *Wnt* level with cell type specification involves the transcription factor *Ascl2*, which interprets a *Wnt* gradient and specifies stem cells by a bimodal switch (Schuijers et al., 2015).

Histone and DNA epigenetic modifications are also important regulators of epithelial stem cell differentiation. Changes in DNA methylation state are associated with the differentiation of hair follicle stem cells toward multiple lineages (Bock et al., 2012). Similarly, DNA methylation dynamics at the enhancers rather than in the proximity of transcription start sites are detected during small intestine stem cell differentiation (Kaaij et al., 2013). A network of epigenetic factors including *Ing5*, *Bptf*, *Smarca5*, *Ezh2*, and *Uhrf1* controls epidermal stem cell fate via a mechanism that includes the regulation of genes that encode proteins responsible for anchoring epidermal stem cells to the extracellular matrix (Mulder et al., 2012). *Cbx4*, a Polycomb Repressive Complex (PRC) 1-associated protein, maintains epidermal stem cells in a slow-cycling and undifferentiated state and protects them from senescence (Luis et al., 2011). In addition, changes in chromatin dynamics allow hair follicle stem cells to cyclically change between quiescent and activated states without losing their stem cell identity (Lien et al., 2011). In the intestine, the mechanisms that allow changes in cell identity between stem cells and differentiated cells involve a permissive chromatin state (Kim et al., 2014).

Chromatin factors can differentially regulate distinct cell compartments within the same epithelium. In adult epidermis *Jarid2* depletion results in delayed hair follicle cycling due to decreased stem cell proliferation, but this does not affect the stem cells of the interfollicular epidermis (Mejetta et al., 2011). Transcription and chromatin factors can locally cooperate to establish gene expression patterns that control stem cell fate. The transcription factor *ZNF750*, for example, interacts with multiple chromatin regulators to repress epidermal stem cell genes and induce differentiation (Boxer et al., 2014). The functional interaction of c-Myc and the histone methyltransferase *Setd8* is required to maintain the sebaceous glands and interfollicular epidermis (Driskell et al., 2012).

The effects of some TFs and chromatin factors only become evident when epithelial homeostasis is disturbed. For example, ablation of the transcriptional co-activator YAP does not affect homeostasis in the intestinal epithelium, yet it is required for



**Figure 7. Transcriptional Noise as an Epithelial Stem Cell Fate Regulator**

Stochastic variations in the expression levels of a single gene or several co-expressed genes are represented as a mechanism to control cell fate decisions. Variation in the amplitude of transcriptional noise within a set of genes is represented as an additional mechanism influencing stem cell fate.

intestinal epithelial repair after injury (Cai et al., 2010). In mammary glands *Ovo12* represses the EMT transcriptional program and is an essential player during tissue regeneration (Watanabe et al., 2014). In skin the PRC2 complex and its repressive histone marks are reduced during wound healing (Shaw and Martin, 2009), consistent with the role of the H3K27 methyltransferases *Ezh2* and *Ezh1* in wound repair (Ezhkova et al., 2009). These examples clearly show the importance of TFs and chromatin factors during tissue damage. However it remains to be uncovered how these factors contribute to epithelial cell plasticity (Figure 5) and how super-enhancer dynamics are involved (Adam et al., 2015).

In summary, the analysis of key players in gene regulation has shown the existence of epigenetic and transcriptional regulators that balance the heterogeneity within an epithelium in terms of differentiation state of cells and lineage identity within a specific epithelial compartment.

### Understanding Cellular Heterogeneity at Single-Cell Resolution

One intriguing aspect of gene transcription is transcriptional noise, which is the variability in gene expression occurring between cells in the same population. Transcriptional noise can control cell fate in hematopoietic lineage selection (Chang et al., 2008), and the noise resulting from transcriptional fluctuations of the pluripotency genes *Sox2*, *Oct4*, and *Nanog* may maintain pluripotency by counteracting differentiation stimuli (Kalmar et al., 2009). Although analysis of cell populations has led to important advances in our knowledge of epithelial stem cell biology, single-cell resolution is required to understand whether phenomena such as transcriptional noise affect epithelial homeostasis and plasticity (Figure 7).

One of the first attempts to understand gene expression by global gene expression profiling in individual epithelial cells iden-

tified *Lrig1* as a stem cell marker and showed that it regulated stem cell quiescence (Jensen and Watt, 2006). Subsequent studies have indicated the existence of two different states of human epidermal stem cells that have not been detected by conventional approaches. Nevertheless, whether they represent distinct stem cell compartments or transitional, interconvertible cell states remains to be explored (Tan et al., 2013).

Single-cell transcriptome analysis in intestinal epithelial cells has shown that the widely used reporters consisting of CreER and EGFP driven by the *Lgr5*, *Bmi1*, and *Hopx* promoters often mark heterogeneous populations and, in addition, that there are discrepancies between reporter activity and endogenous transcripts. This analysis has also highlighted that *Lgr5*, *Bmi1*, and *Hopx* cell populations are molecularly and functionally distinct, supporting a role for two distinct stem cell populations in intestinal maintenance (Li et al., 2014). Single-cell genomic approaches can provide an understanding of stem cell dynamics at higher resolution than by lineage tracing alone, as illustrated recently for lung epithelium (Treutlein et al., 2014). In the future, combining single-cell (proteomic and genomic) approaches with lineage tracing will allow cross-validation of the conclusions and further strengthen our understanding of epithelial stem cells.

Analysis of single-cell data has been greatly enriched by the development of new computational methods. For instance by applying the Wanderlust algorithm to single-cell mass cytometry data, it was possible to construct trajectories of how hematopoietic stem cells differentiate into B cells, identifying new cell states and aligning them to specific signaling pathways (Bendall et al., 2014). To robustly identify previously hidden cell subpopulations, computational methods can eliminate the effects of potential confounding factors (such as cell cycle stage) on the heterogeneity of gene expression identified by single-cell RNA sequencing (Buettner et al., 2015). Single-cell RNA-seq data have also been analyzed using the self-organizing map (SOM), which represents a way to visualize and interrogate cell-to-cell heterogeneity based on the behavior of coordinately expressed gene clusters (Kim et al., 2015).

The high-throughput data provided by RNA-seq experiments lacks positional information. This is not a problem with blood cells in the circulation, but it is a major disadvantage for analyzing epithelial cells, since spatial organization often reflects cellular heterogeneity. In order to overcome this problem, a recent method called FISSEQ (fluorescence in situ RNA sequencing) has been developed that combines single-cell transcriptome analysis with a spatial context (Lee et al., 2014).

### Generation of Epithelial Tissues from Pluripotent Stem Cells

Techniques for regenerating epidermis in vitro date back many decades, but progress in regenerating other epithelia in culture is more recent. Important advances include techniques for growing epithelial organoids from single cells and recapitulating the differentiated lineages that are present in the normal tissue. It is possible to grow intestinal organoids from purified LGR5-positive stem cells such that a single crypt can generate villus-like epithelial domains in which all differentiated cell types are present in the absence of a non-epithelial cellular niche (Sato et al., 2009) (Figure 4B). Similar culture conditions allow growth of epithelial organoids from human small intestine and colon

(Sato et al., 2011), which could be used as a resource for drug screening or to graft genotype-matched human intestine and colon epithelium. A 3D culture system to maintain mouse and human prostate organoids has also been developed (Karthaus et al., 2014). It is also possible to recreate salivary glands and hair follicles in culture (Nanduri et al., 2014; Tobin, 2011).

An extension of these studies is to generate epithelial tissues from human pluripotent stem cells. This offers the possibility of generating not only the epithelia themselves, but also associated cell types, such as melanocytes and fibroblasts in the case of skin (Bilousova and Roop, 2014). Progress is not restricted to skin, as gastric organoids have been successfully generated from pluripotent cells (McCracken et al., 2014). It is yet to be determined how similar the epithelia generated from pluripotent cells are to adult or fetal tissues, as immaturity is a concern for other differentiated cells types, such as liver (Stutchfield et al., 2010).

### Looking Forward

In recent years we have gained unprecedented insights into the nature and regulation of stem cells in a variety of postnatal epithelia, finding common principles such as stem cell compartmentalization and plasticity. Looking to the future, single-cell genome-wide technologies will give us new perspectives on cell identity and the interplay between epigenetic and transcriptional regulation. The creation of entire epithelial organs from pluripotent stem cells will have practical applications in disease modeling and drug discovery, while modulating the stem cell niche via changes in stromal cells, immune cells, and bacteria will improve the quality of tissue repair for regenerative medicine purposes.

### ACKNOWLEDGMENTS

We gratefully acknowledge input from all our colleagues (<http://www.wattlab.org>). We thank the Wellcome Trust, Medical Research Council, BBSRC, and UKRMP for funding.

### REFERENCES

Adam, R.C., Yang, H., Rockowitz, S., Larsen, S.B., Nikolova, M., Oristian, D.S., Polak, L., Kadaja, M., Asare, A., Zheng, D., and Fuchs, E. (2015). Pioneer factors govern super-enhancer dynamics in stem cell plasticity and lineage choice. *Nature*, in press. Published online March 18, 2015. <http://dx.doi.org/10.1038/nature14289>.

Alcolea, M.P., Greulich, P., Wabik, A., Frede, J., Simons, B.D., and Jones, P.H. (2014). Differentiation imbalance in single oesophageal progenitor cells causes clonal immortalization and field change. *Nat. Cell Biol.* **16**, 615–622.

Anderson, G., and Takahama, Y. (2012). Thymic epithelial cells: working class heroes for T cell development and repertoire selection. *Trends Immunol.* **33**, 256–263.

Baker, A.M., Cereser, B., Melton, S., Fletcher, A.G., Rodriguez-Justo, M., Tardous, P.J., Humphries, A., Elia, G., McDonald, S.A., Wright, N.A., et al. (2014). Quantification of crypt and stem cell evolution in the normal and neoplastic human colon. *Cell Rep.* **8**, 940–947.

Barker, N., van Es, J.H., Kuipers, J., Kujala, P., van den Born, M., Cozijnsen, M., Haegebarth, A., Korving, J., Begthel, H., Peters, P.J., and Clevers, H. (2007). Identification of stem cells in small intestine and colon by marker gene Lgr5. *Nature* **449**, 1003–1007.

Bendall, S.C., Davis, K.L., Amir, A.D., Tadmor, M.D., Simonds, E.F., Chen, T.J., Shenfeld, D.K., Nolan, G.P., and Pe'er, D. (2014). Single-cell trajectory detection uncovers progression and regulatory coordination in human B cell development. *Cell* **157**, 714–725.

Bilousova, G., and Roop, D.R. (2014). Induced pluripotent stem cells in dermatology: potentials, advances, and limitations. *Cold Spring Harbor Perspect. Med.* **4**, a015164.

Blanpain, C., and Simons, B.D. (2013). Unravelling stem cell dynamics by lineage tracing. *Nat. Rev. Mol. Cell Biol.* **14**, 489–502.

Bock, C., Beerman, I., Lien, W.H., Smith, Z.D., Gu, H., Boyle, P., Gnirke, A., Fuchs, E., Rossi, D.J., and Meissner, A. (2012). DNA methylation dynamics during in vivo differentiation of blood and skin stem cells. *Mol. Cell* **47**, 633–647.

Boxer, L.D., Barajas, B., Tao, S., Zhang, J., and Khavari, P.A. (2014). ZNF750 interacts with KLF4 and RCOR1, KDM1A, and CTBP1/2 chromatin regulators to repress epidermal progenitor genes and induce differentiation genes. *Genes Dev.* **28**, 2013–2026.

Buczacki, S.J., Zecchini, H.I., Nicholson, A.M., Russell, R., Vermeulen, L., Kemp, R., and Winton, D.J. (2013). Intestinal label-retaining cells are secretory precursors expressing Lgr5. *Nature* **495**, 65–69.

Buettner, F., Natarajan, K.N., Casale, F.P., Proserpio, V., Scialdone, A., Theis, F.J., Teichmann, S.A., Marioni, J.C., and Stegle, O. (2015). Computational analysis of cell-to-cell heterogeneity in single-cell RNA-sequencing data reveals hidden subpopulations of cells. *Nat. Biotechnol.* **33**, 155–160.

Busch, K., Klapproth, K., Barile, M., Flossdorf, M., Holland-Letz, T., Schlenner, S.M., Reth, M., Höfer, T., and Rodewald, H.R. (2015). Fundamental properties of unperturbed haematopoiesis from stem cells in vivo. *Nature* **518**, 542–546.

Cai, J., Zhang, N., Zheng, Y., de Wilde, R.F., Maitra, A., and Pan, D. (2010). The Hippo signaling pathway restricts the oncogenic potential of an intestinal regeneration program. *Genes Dev.* **24**, 2383–2388.

Calvo, F., Ege, N., Grande-Garcia, A., Hooper, S., Jenkins, R.P., Chaudhry, S.I., Harrington, K., Williamson, P., Moeendarbary, E., Charras, G., and Sahai, E. (2013). Mechanotransduction and YAP-dependent matrix remodeling is required for the generation and maintenance of cancer-associated fibroblasts. *Nat. Cell Biol.* **15**, 637–646.

Chang, H.H., Hemberg, M., Barahona, M., Ingber, D.E., and Huang, S. (2008). Transcriptome-wide noise controls lineage choice in mammalian progenitor cells. *Nature* **453**, 544–547.

Clayton, E., Doupé, D.P., Klein, A.M., Winton, D.J., Simons, B.D., and Jones, P.H. (2007). A single type of progenitor cell maintains normal epidermis. *Nature* **446**, 185–189.

DeWard, A.D., Cramer, J., and Lagasse, E. (2014). Cellular heterogeneity in the mouse esophagus implicates the presence of a nonquiescent epithelial stem cell population. *Cell Rep.* **9**, 701–711.

Donati, G., Proserpio, V., Lichtenberger, B.M., Natsuga, K., Sinclair, R., Fujiwara, H., and Watt, F.M. (2014). Epidermal Wnt/ $\beta$ -catenin signaling regulates adipocyte differentiation via secretion of adipogenic factors. *Proc. Natl. Acad. Sci. USA* **111**, E1501–E1509.

Driessens, G., Beck, B., Caauwe, A., Simons, B.D., and Blanpain, C. (2012). Defining the mode of tumour growth by clonal analysis. *Nature* **488**, 527–530.

Driskell, I., Oda, H., Blanco, S., Nascimento, E., Humphreys, P., and Frye, M. (2012). The histone methyltransferase Setd8 acts in concert with c-Myc and is required to maintain skin. *EMBO J.* **31**, 616–629.

Driskell, R.R., Lichtenberger, B.M., Hoste, E., Kretschmar, K., Simons, B.D., Charalambous, M., Ferron, S.R., Herault, Y., Pavlovic, G., Ferguson-Smith, A.C., and Watt, F.M. (2013). Distinct fibroblast lineages determine dermal architecture in skin development and repair. *Nature* **504**, 277–281.

Estrach, S., Cordes, R., Hozumi, K., Gossler, A., and Watt, F.M. (2008). Role of the Notch ligand Delta1 in embryonic and adult mouse epidermis. *J. Invest. Dermatol.* **128**, 825–832.

Ezhkova, E., Pasolli, H.A., Parker, J.S., Stokes, N., Su, I.H., Hannon, G., Tarakhovskiy, A., and Fuchs, E. (2009). Ezh2 orchestrates gene expression for the stepwise differentiation of tissue-specific stem cells. *Cell* **136**, 1122–1135.

Festa, E., Fretz, J., Berry, R., Schmidt, B., Rodeheffer, M., Horowitz, M., and Horsley, V. (2011). Adipocyte lineage cells contribute to the skin stem cell niche to drive hair cycling. *Cell* **146**, 761–771.

- Folgueras, A.R., Guo, X., Pasolli, H.A., Stokes, N., Polak, L., Zheng, D., and Fuchs, E. (2013). Architectural niche organization by LHX2 is linked to hair follicle stem cell function. *Cell Stem Cell* **13**, 314–327.
- Fujiwara, H., Ferreira, M., Donati, G., Marciano, D.K., Linton, J.M., Sato, Y., Hartner, A., Sekiguchi, K., Reichardt, L.F., and Watt, F.M. (2011). The basement membrane of hair follicle stem cells is a muscle cell niche. *Cell* **144**, 577–589.
- Gay, D., Kwon, O., Zhang, Z., Spata, M., Plikus, M.V., Holler, P.D., Ito, M., Yang, Z., Treffeisen, E., Kim, C.D., et al. (2013). Fgf9 from dermal  $\gamma\delta$  T cells induces hair follicle neogenesis after wounding. *Nat. Med.* **19**, 916–923.
- Gollwitzer, E.S., Saglani, S., Trompette, A., Yadava, K., Sherburn, R., McCoy, K.D., Nicod, L.P., Lloyd, C.M., and Marsland, B.J. (2014). Lung microbiota promotes tolerance to allergens in neonates via PD-L1. *Nat. Med.* **20**, 642–647.
- Gomez, C., Chua, W., Miremadi, A., Quist, S., Headon, D.J., and Watt, F.M. (2013). The interfollicular epidermis of adult mouse tail comprises two distinct cell lineages that are differentially regulated by Wnt, Edaradd, and Lrig1. *Stem Cell Reports* **1**, 19–27.
- Grice, E.A., and Segre, J.A. (2011). The skin microbiome. *Nat. Rev. Microbiol.* **9**, 244–253.
- Grompe, M. (2014). Liver stem cells, where art thou? *Cell Stem Cell* **15**, 257–258.
- Hall, P.A., and Watt, F.M. (1989). Stem cells: the generation and maintenance of cellular diversity. *Development* **106**, 619–633.
- Hnisz, D., Abraham, B.J., Lee, T.I., Lau, A., Saint-André, V., Sigova, A.A., Hoke, H.A., and Young, R.A. (2013). Super-enhancers in the control of cell identity and disease. *Cell* **155**, 934–947.
- Hogan, B.L., Barkauskas, C.E., Chapman, H.A., Epstein, J.A., Jain, R., Hsia, C.C., Niklason, L., Calle, E., Le, A., Randell, S.H., et al. (2014). Repair and regeneration of the respiratory system: complexity, plasticity, and mechanisms of lung stem cell function. *Cell Stem Cell* **15**, 123–138.
- Hoste, E., Arwert, E.N., Lal, R., South, A.P., Salas-Alanis, J.C., Murrell, D.F., Donati, G., and Watt, F.M. (2015). Innate sensing of microbial products promotes wound-induced skin cancer. *Nat. Commun.* **6**, 5932.
- Hsu, Y.C., Li, L., and Fuchs, E. (2014). Transit-amplifying cells orchestrate stem cell activity and tissue regeneration. *Cell* **157**, 935–949.
- Hu, B., Castillo, E., Harewood, L., Ostano, P., Raymond, A., Dummer, R., Raf-foul, W., Hoetzenecker, W., Hofbauer, G.F., and Dotto, G.P. (2012). Multifocal epithelial tumors and field cancerization from loss of mesenchymal CSL signaling. *Cell* **149**, 1207–1220.
- Ito, M., Liu, Y., Yang, Z., Nguyen, J., Liang, F., Morris, R.J., and Cotsarelis, G. (2005). Stem cells in the hair follicle bulge contribute to wound repair but not to homeostasis of the epidermis. *Nat. Med.* **11**, 1351–1354.
- Ito, M., Yang, Z., Andl, T., Cui, C., Kim, N., Millar, S.E., and Cotsarelis, G. (2007). Wnt-dependent de novo hair follicle regeneration in adult mouse skin after wounding. *Nature* **447**, 316–320.
- Izkovitz, S., Lyubimova, A., Blat, I.C., Maynard, M., van Es, J., Lees, J., Jacks, T., Clevers, H., and van Oudenaarden, A. (2012). Single-molecule transcript counting of stem-cell markers in the mouse intestine. *Nat. Cell Biol.* **14**, 106–114.
- Jaks, V., Barker, N., Kasper, M., van Es, J.H., Snippert, H.J., Clevers, H., and Toftgård, R. (2008). Lgr5 marks cycling, yet long-lived, hair follicle stem cells. *Nat. Genet.* **40**, 1291–1299.
- Jensen, K.B., and Watt, F.M. (2006). Single-cell expression profiling of human epidermal stem and transit-amplifying cells: Lrig1 is a regulator of stem cell quiescence. *Proc. Natl. Acad. Sci. USA* **103**, 11958–11963.
- Jensen, K.B., Collins, C.A., Nascimento, E., Tan, D.W., Frye, M., Itami, S., and Watt, F.M. (2009). Lrig1 expression defines a distinct multipotent stem cell population in mammalian epidermis. *Cell Stem Cell* **4**, 427–439.
- Kaaij, L.T., van de Wetering, M., Fang, F., Decato, B., Molaro, A., van de Werken, H.J., van Es, J.H., Schuijers, J., de Wit, E., de Laat, W., et al. (2013). DNA methylation dynamics during intestinal stem cell differentiation reveals enhancers driving gene expression in the villus. *Genome Biol.* **14**, R50.
- Kalmar, T., Lim, C., Hayward, P., Muñoz-Descalzo, S., Nichols, J., Garcia-Ojalvo, J., and Martinez Arias, A. (2009). Regulated fluctuations in nanog expression mediate cell fate decisions in embryonic stem cells. *PLoS Biol.* **7**, e1000149.
- Karthaus, W.R., Iaquina, P.J., Drost, J., Gracanin, A., van Boxtel, R., Wongvipat, J., Dowling, C.M., Gao, D., Begthel, H., Sachs, N., et al. (2014). Identification of multipotent luminal progenitor cells in human prostate organoid cultures. *Cell* **159**, 163–175.
- Kaukua, N., Shahidi, M.K., Konstantinidou, C., Dyachuk, V., Kaucska, M., Furlan, A., An, Z., Wang, L., Hultman, I., Ahrlund-Richter, L., et al. (2014). Glial origin of mesenchymal stem cells in a tooth model system. *Nature* **513**, 551–554.
- Kim, T.H., Li, F., Ferreiro-Neira, I., Ho, L.L., Luyten, A., Nalapareddy, K., Long, H., Verzi, M., and Shvadasani, R.A. (2014). Broadly permissive intestinal chromatin underlies lateral inhibition and cell plasticity. *Nature* **506**, 511–515.
- Kim, D.H., Marinov, G.K., Pepke, S., Singer, Z.S., He, P., Williams, B., Schroth, G.P., Elowitz, M.B., and Wold, B.J. (2015). Single-cell transcriptome analysis reveals dynamic changes in lncRNA expression during reprogramming. *Cell Stem Cell* **16**, 88–101.
- Klein, A.M., Brash, D.E., Jones, P.H., and Simons, B.D. (2010). Stochastic fate of p53-mutant epidermal progenitor cells is tilted toward proliferation by UV B during preneoplasia. *Proc. Natl. Acad. Sci. USA* **107**, 270–275.
- Kretzschmar, K., and Watt, F.M. (2012). Lineage tracing. *Cell* **148**, 33–45.
- Kumar, M.E., Bogard, P.E., Espinoza, F.H., Menke, D.B., Kingsley, D.M., and Krasnow, M.A. (2014). Mesenchymal cells. Defining a mesenchymal progenitor niche at single-cell resolution. *Science* **346**, 1258810.
- Lane, S.W., Williams, D.A., and Watt, F.M. (2014). Modulating the stem cell niche for tissue regeneration. *Nature biotechnology* **32**, 795–803.
- Lee, J.H., Daugharthy, E.R., Scheiman, J., Kalhor, R., Yang, J.L., Ferrante, T.C., Terry, R., Jeanty, S.S., Li, C., Amamoto, R., et al. (2014). Highly multiplexed subcellular RNA sequencing in situ. *Science* **343**, 1360–1363.
- Li, L., and Clevers, H. (2010). Coexistence of quiescent and active adult stem cells in mammals. *Science* **327**, 542–545.
- Li, W., Qiao, W., Chen, L., Xu, X., Yang, X., Li, D., Li, C., Brodie, S.G., Meguid, M.M., Hennighausen, L., and Deng, C.X. (2003). Squamous cell carcinoma and mammary abscess formation through squamous metaplasia in Smad4/Dpc4 conditional knockout mice. *Development* **130**, 6143–6153.
- Li, N., Yousefi, M., Nakauka-Ddamba, A., Jain, R., Tobias, J., Epstein, J.A., Jensen, S.T., and Lengner, C.J. (2014). Single-cell analysis of proxy reporter allele-marked epithelial cells establishes intestinal stem cell hierarchy. *Stem Cell Reports* **3**, 876–891.
- Lien, W.H., Guo, X., Polak, L., Lawton, L.N., Young, R.A., Zheng, D., and Fuchs, E. (2011). Genome-wide maps of histone modifications unwind in vivo chromatin states of the hair follicle lineage. *Cell Stem Cell* **9**, 219–232.
- Lim, X., Tan, S.H., Koh, W.L., Chau, R.M., Yan, K.S., Kuo, C.J., van Amerongen, R., Klein, A.M., and Nusse, R. (2013). Interfollicular epidermal stem cells self-renew via autocrine Wnt signaling. *Science* **342**, 1226–1230.
- Luis, N.M., Morey, L., Mejetta, S., Pascual, G., Janich, P., Kuebler, B., Cozutto, L., Roma, G., Nascimento, E., Frye, M., et al. (2011). Regulation of human epidermal stem cell proliferation and senescence requires polycomb-dependent and -independent functions of Cbx4. *Cell Stem Cell* **9**, 233–246.
- Mascre, G., Dekoninck, S., Drogat, B., Youssef, K.K., Broheé, S., Sotiropoulou, P.A., Simons, B.D., and Blanpain, C. (2012). Distinct contribution of stem and progenitor cells to epidermal maintenance. *Nature* **489**, 257–262.
- McCracken, K.W., Catá, E.M., Crawford, C.M., Sinagoga, K.L., Schumacher, M., Rockich, B.E., Tsai, Y.H., Mayhew, C.N., Spence, J.R., Zavros, Y., and Wells, J.M. (2014). Modelling human development and disease in pluripotent stem-cell-derived gastric organoids. *Nature* **516**, 400–404.
- Mejetta, S., Morey, L., Pascual, G., Kuebler, B., Mysliwiec, M.R., Lee, Y., Shiekh-tarrar, R., Di Croce, L., and Benitah, S.A. (2011). Jarid2 regulates mouse epidermal stem cell activation and differentiation. *EMBO J.* **30**, 3635–3646.
- Mouw, J.K., Yui, Y., Damiano, L., Bainer, R.O., Lakins, J.N., Acerbi, I., Ou, G., Wijekoon, A.C., Levental, K.R., Gilbert, P.M., et al. (2014). Tissue mechanics

modulate microRNA-dependent PTEN expression to regulate malignant progression. *Nat. Med.* **20**, 360–367.

Mulder, K.W., Wang, X., Escriu, C., Ito, Y., Schwarz, R.F., Gillis, J., Sirokmány, G., Donati, G., Uribe-Lewis, S., Pavlidis, P., et al. (2012). Diverse epigenetic strategies interact to control epidermal differentiation. *Nat. Cell Biol.* **14**, 753–763.

Muñoz, J., Stange, D.E., Schepers, A.G., van de Wetering, M., Koo, B.K., Itzkovitz, S., Volckmann, R., Kung, K.S., Koster, J., Radulescu, S., et al. (2012). The Lgr5 intestinal stem cell signature: robust expression of proposed quiescent '4+4' cell markers. *EMBO J.* **31**, 3079–3091.

Naik, S., Bouladoux, N., Linehan, J.L., Han, S.J., Harrison, O.J., Wilhelm, C., Conlan, S., Himmelfarb, S., Byrd, A.L., Deming, C., et al. (2015). Commensal-dendritic-cell interaction specifies a unique protective skin immune signature. *Nature* **520**, 104–108.

Nanduri, L.S., Baanstra, M., Faber, H., Rocchi, C., Zwart, E., de Haan, G., van Os, R., and Coppes, R.P. (2014). Purification and ex vivo expansion of fully functional salivary gland stem cells. *Stem Cell Reports* **3**, 957–964.

Nigro, G., Rossi, R., Commere, P.H., Jay, P., and Sansonetti, P.J. (2014). The cytosolic bacterial peptidoglycan sensor Nod2 affords stem cell protection and links microbes to gut epithelial regeneration. *Cell Host Microbe* **15**, 792–798.

Öhlund, D., Elyada, E., and Tuveson, D. (2014). Fibroblast heterogeneity in the cancer wound. *J. Exp. Med.* **211**, 1503–1523.

Page, M.E., Lombard, P., Ng, F., Göttgens, B., and Jensen, K.B. (2013). The epidermis comprises autonomous compartments maintained by distinct stem cell populations. *Cell Stem Cell* **13**, 471–482.

Pasparakis, M., Haase, I., and Nestle, F.O. (2014). Mechanisms regulating skin immunity and inflammation. *Nat. Rev. Immunol.* **14**, 289–301.

Powell, A.E., Wang, Y., Li, Y., Poulin, E.J., Means, A.L., Washington, M.K., Higginbotham, J.N., Juchheim, A., Prasad, N., Levy, S.E., et al. (2012). The pan-ErbB negative regulator Lrig1 is an intestinal stem cell marker that functions as a tumor suppressor. *Cell* **149**, 146–158.

Prater, M.D., Petit, V., Alasdair Russell, I., Girardi, R.R., Shehata, M., Menon, S., Schulte, R., Kalajzic, I., Rath, N., Olson, M.F., et al. (2014). Mammary stem cells have myoepithelial cell properties. *Nat. Cell Biol.* **16**, 942–950.

Rawlins, E.L., Okubo, T., Xue, Y., Brass, D.M., Auten, R.L., Hasegawa, H., Wang, F., and Hogan, B.L. (2009). The role of Scgb1a1+ Clara cells in the long-term maintenance and repair of lung airway, but not alveolar, epithelium. *Cell Stem Cell* **4**, 525–534.

Rinkevich, Y., Lindau, P., Ueno, H., Longaker, M.T., and Weissman, I.L. (2011). Germ-layer and lineage-restricted stem/progenitors regenerate the mouse digit tip. *Nature* **476**, 409–413.

Rinkevich, Y., Walmsley, G.G., Hu, M.S., Maan, Z.N., Newman, A.M., Drukker, M., Januszzyk, M., Krampitz, G.W., Gurtner, G.C., Lorenz, H.P., et al. (2015). Skin fibrosis. Identification and isolation of a dermal lineage with intrinsic fibrogenic potential. *Science* **348**, aaa2151.

Rios, A.C., Fu, N.Y., Lindeman, G.J., and Visvader, J.E. (2014). In situ identification of bipotent stem cells in the mammary gland. *Nature* **506**, 322–327.

Ritsma, L., Ellenbroek, S.I., Zomer, A., Snippert, H.J., de Sauvage, F.J., Simons, B.D., Clevers, H., and van Rheenen, J. (2014). Intestinal crypt homeostasis revealed at single-stem-cell level by in vivo live imaging. *Nature* **507**, 362–365.

Rompolas, P., Mesa, K.R., and Greco, V. (2013). Spatial organization within a niche as a determinant of stem-cell fate. *Nature* **502**, 513–518.

Rossi, D.J., Jamieson, C.H., and Weissman, I.L. (2008). Stems cells and the pathways to aging and cancer. *Cell* **132**, 681–696.

Sangiorgi, E., and Capecchi, M.R. (2008). Bmi1 is expressed in vivo in intestinal stem cells. *Nat. Genet.* **40**, 915–920.

Sato, T., Vries, R.G., Snippert, H.J., van de Wetering, M., Barker, N., Stange, D.E., van Es, J.H., Abo, A., Kujala, P., Peters, P.J., and Clevers, H. (2009). Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature* **459**, 262–265.

Sato, T., Stange, D.E., Ferrante, M., Vries, R.G., Van Es, J.H., Van den Brink, S., Van Houdt, W.J., Pronk, A., Van Gorp, J., Siersema, P.D., and Clevers, H. (2011). Long-term expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and Barrett's epithelium. *Gastroenterology* **141**, 1762–1772.

Schaub, J.R., Malato, Y., Gormond, C., and Willenbring, H. (2014). Evidence against a stem cell origin of new hepatocytes in a common mouse model of chronic liver injury. *Cell Rep.* **8**, 933–939.

Schepers, A.G., Snippert, H.J., Stange, D.E., van den Born, M., van Es, J.H., van de Wetering, M., and Clevers, H. (2012). Lineage tracing reveals Lgr5+ stem cell activity in mouse intestinal adenomas. *Science* **337**, 730–735.

Scherz-Shouval, R., Santagata, S., Mendillo, M.L., Sholl, L.M., Ben-Aharon, I., Beck, A.H., Dias-Santagata, D., Koeva, M., Stemmer, S.M., Whitesell, L., and Lindquist, S. (2014). The reprogramming of tumor stroma by HSF1 is a potent enabler of malignancy. *Cell* **158**, 564–578.

Schuijers, J., Junker, J.P., Mokry, M., Hatzis, P., Koo, B.K., Sasselli, V., van der Flier, L.G., Cuppen, E., van Oudenaarden, A., and Clevers, H. (2015). *Ascl2* acts as an R-spondin/Wnt-responsive switch to control stemness in intestinal crypts. *Cell Stem Cell* **16**, 158–170.

Shaw, T., and Martin, P. (2009). Epigenetic reprogramming during wound healing: loss of polycomb-mediated silencing may enable upregulation of repair genes. *EMBO Rep.* **10**, 881–886.

Sigal, M., Rothenberg, M.E., Logan, C.Y., Lee, J.Y., Honaker, R.W., Cooper, R.L., Passarelli, B., Camorlinga, M., Bouley, D.M., Alvarez, G., et al. (2015). *Helicobacter pylori* Activates and Expands Lgr5 Stem Cells Through Direct Colonization of the Gastric Glands. *Gastroenterology*, in press. Published online February 26, 2015. <http://dx.doi.org/10.1053/j.gastro.2015.02.049>.

Simmini, S., Bialecka, M., Huch, M., Kester, L., van de Wetering, M., Sato, T., Beck, F., van Oudenaarden, A., Clevers, H., and Deschamps, J. (2014). Trans-formation of intestinal stem cells into gastric stem cells on loss of transcription factor Cdx2. *Nat. Commun.* **5**, 5728.

Slack, J.M. (2007). Metaplasia and transdifferentiation: from pure biology to the clinic. *Nat. Rev. Mol. Cell Biol.* **8**, 369–378.

Snippert, H.J., van der Flier, L.G., Sato, T., van Es, J.H., van den Born, M., Kroon-Veenboer, C., Barker, N., Klein, A.M., van Rheenen, J., Simons, B.D., and Clevers, H. (2010). Intestinal crypt homeostasis results from neutral competition between symmetrically dividing Lgr5 stem cells. *Cell* **143**, 134–144.

Snippert, H.J., Schepers, A.G., van Es, J.H., Simons, B.D., and Clevers, H. (2014). Biased competition between Lgr5 intestinal stem cells driven by oncogenic mutation induces clonal expansion. *EMBO Rep.* **15**, 62–69.

Stutchfield, B.M., Forbes, S.J., and Wigmore, S.J. (2010). Prospects for stem cell transplantation in the treatment of hepatic disease. *Liver Transpl.* **16**, 827–836.

Sugimoto, H., Mundel, T.M., Kieran, M.W., and Kalluri, R. (2006). Identification of fibroblast heterogeneity in the tumor microenvironment. *Cancer Biol. Ther.* **5**, 1640–1646.

Sun, J., Ramos, A., Chapman, B., Johnnidis, J.B., Le, L., Ho, Y.J., Klein, A., Hofmann, O., and Camargo, F.D. (2014). Clonal dynamics of native haematopoiesis. *Nature* **514**, 322–327.

Takeda, N., Jain, R., LeBoeuf, M.R., Wang, Q., Lu, M.M., and Epstein, J.A. (2011). Interconversion between intestinal stem cell populations in distinct niches. *Science* **334**, 1420–1424.

Takeo, M., Chou, W.C., Sun, Q., Lee, W., Rabbani, P., Loomis, C., Taketo, M.M., and Ito, M. (2013). Wnt activation in nail epithelium couples nail growth to digit regeneration. *Nature* **499**, 228–232.

Tan, D.W., Jensen, K.B., Trotter, M.W., Connelly, J.T., Broad, S., and Watt, F.M. (2013). Single-cell gene expression profiling reveals functional heterogeneity of undifferentiated human epidermal cells. *Development* **140**, 1433–1444.

Tarlow, B.D., Pelz, C., Naugler, W.E., Wakefield, L., Wilson, E.M., Finegold, M.J., and Grompe, M. (2014). Bipotent adult liver progenitors are derived from chronically injured mature hepatocytes. *Cell Stem Cell* **15**, 605–618.

- Tata, P.R., Mou, H., Pardo-Saganta, A., Zhao, R., Prabhu, M., Law, B.M., Vinarsky, V., Cho, J.L., Breton, S., Sahay, A., et al. (2013). Dedifferentiation of committed epithelial cells into stem cells in vivo. *Nature* 503, 218–223.
- Thorel, F., Népote, V., Avril, I., Kohno, K., Desgraz, R., Chera, S., and Herrera, P.L. (2010). Conversion of adult pancreatic alpha-cells to beta-cells after extreme beta-cell loss. *Nature* 464, 1149–1154.
- Tian, H., Biehs, B., Warming, S., Leong, K.G., Rangell, L., Klein, O.D., and de Sauvage, F.J. (2011). A reserve stem cell population in small intestine renders Lgr5-positive cells dispensable. *Nature* 478, 255–259.
- Tobin, D.J. (2011). Ex vivo organ culture of human hair follicles: a model epithelial-neuroectodermal-mesenchymal interaction system. *Methods Mol. Biol.* 695, 213–227.
- Treutlein, B., Brownfield, D.G., Wu, A.R., Neff, N.F., Mantalas, G.L., Espinoza, F.H., Desai, T.J., Krasnow, M.A., and Quake, S.R. (2014). Reconstructing lineage hierarchies of the distal lung epithelium using single-cell RNA-seq. *Nature* 509, 371–375.
- van Es, J.H., Sato, T., van de Wetering, M., Lyubimova, A., Nee, A.N., Gregorieff, A., Sasaki, N., Zeinstra, L., van den Born, M., Korving, J., et al. (2012). Dll1+ secretory progenitor cells revert to stem cells upon crypt damage. *Nat. Cell Biol.* 14, 1099–1104.
- Van Keymeulen, A., Rocha, A.S., Ousset, M., Beck, B., Bouvencourt, G., Rock, J., Sharma, N., Dekoninck, S., and Blanpain, C. (2011). Distinct stem cells contribute to mammary gland development and maintenance. *Nature* 479, 189–193.
- Vermeulen, L., Morrissey, E., van der Heijden, M., Nicholson, A.M., Sottoriva, A., Buczacck, S., Kemp, R., Tavaré, S., and Winton, D.J. (2013). Defining stem cell dynamics in models of intestinal tumor initiation. *Science* 342, 995–998.
- Visvader, J.E., and Stingl, J. (2014). Mammary stem cells and the differentiation hierarchy: current status and perspectives. *Genes Dev.* 28, 1143–1158.
- Watanabe, K., Villarreal-Ponce, A., Sun, P., Salmans, M.L., Fallahi, M., Andersen, B., and Dai, X. (2014). Mammary morphogenesis and regeneration require the inhibition of EMT at terminal end buds by *Ovol2* transcriptional repressor. *Dev. Cell* 29, 59–74.
- Watt, F.M. (2014). Mammalian skin cell biology: at the interface between laboratory and clinic. *Science* 346, 937–940.
- Watt, F.M., and Fujiwara, H. (2011). Cell-extracellular matrix interactions in normal and diseased skin. *Cold Spring Harb. Perspect. Biol.* 3.
- Yanger, K., Zong, Y., Maggs, L.R., Shapira, S.N., Maddipati, R., Aiello, N.M., Thung, S.N., Wells, R.G., Greenbaum, L.E., and Stanger, B.Z. (2013). Robust cellular reprogramming occurs spontaneously during liver regeneration. *Genes Dev.* 27, 719–724.
- Zhang, L.J., Guerrero-Juarez, C.F., Hata, T., Bapat, S.P., Ramos, R., Plikus, M.V., and Gallo, R.L. (2015). Innate immunity. Dermal adipocytes protect against invasive *Staphylococcus aureus* skin infection. *Science* 347, 67–71.
- Zomer, A., Ellenbroek, S.I., Ritsma, L., Beerling, E., Vrisekoop, N., and Van Rheenen, J. (2013). Intravital imaging of cancer stem cell plasticity in mammary tumors. *Stem Cells* 31, 602–606.
- Zuo, W., Zhang, T., Wu, D.Z., Guan, S.P., Liew, A.A., Yamamoto, Y., Wang, X., Lim, S.J., Vincent, M., Lessard, M., et al. (2015). p63(+)Krt5(+) distal airway stem cells are essential for lung regeneration. *Nature* 517, 616–620.