## Preliminary Examination 1998 Complex Analysis

Do all problems.

Notation:

$\mathbb{R} = \{ x : x \text{ is a real number } \}$	$\mathbb{C} = \{ z : z \text{ is a complex number } \}$
$B(a,r) = \{ z \in \mathbb{C} :   z - a   < r \}$	ann $(a, r_1, r_2) = \{ z \in \mathbb{C} : r_1 <   z - a   < r_2 \}$

For  $G \subset \mathbb{C}$ , let  $\mathcal{A}(G)$  denote the set of analytic functions on G (mapping G to  $\mathbb{C}$ ).

- 1. Let f be an entire function.
  - (a) Suppose there exist  $a, b \in \mathbb{R}$  such that  $|f(z)| \leq (a\sqrt{|z|} + b)$  for all  $z \in \mathbb{C}$ . Show that f is constant.
  - (b) Suppose there exist  $a, b \in \mathbb{R}$  such that  $|f(z)| \leq (a |z|^{5/2} + b)$  for all  $z \in \mathbb{C}$ . What can you say about *f*?
- 2. Let *f*, *g* be entire functions. Suppose  $|f(z)| \le |g(z)|$  for all  $z \in \mathbb{C}$ . Prove there exists a constant *c* such that  $f \equiv cg$ .
- 3. Let A(r) = ann(0,r,1), 0 < r < 1, and  $B = B(0,1) \setminus B(\frac{1}{4},\frac{1}{4})$ . Show that there exists an *r* such that A(r) is conformally equivalent to *B*.
- 4. Let  $G_1$  and  $G_2$  be simply connected regions, neither region is all of  $\mathbb{C}$ . Let  $a \in G_1$ . Suppose that  $f, g \in \mathcal{A}(G_1)$  such that f is one-to-one on  $G_1$  with  $f(G_1) = G_2$ ,  $g(G_1) \subset G_2$  and f(a) = g(a). Prove  $|g'(a)| \leq |f'(a)|$ .
- 5. Define the Gamma function,  $\Gamma$ . Prove that  $\Gamma(z)\Gamma(1-z) = \pi/\sin(\pi z)$ , for *z* not an integer.
- 6. Let G be a region in  $\mathbb{C}$  and let  $\mathcal{F}$  be a subset of  $\mathcal{A}(G)$ . Prove that if  $\mathcal{F}$  is locally bounded, then  $\mathcal{F}$  is equicontinuous at each point of G.
- 7. Let  $D_1 = \{ z \in B(0,1) : \text{Im } z > \frac{1}{2} \}$ . Find a conformal map f which maps  $D_1$  one-to-one and onto B(0,1) such that f(3i/4) = 0.

8. Compute 
$$\int_{-\infty}^{\infty} \frac{e^{iax}}{(1+x^2)^2} dx, a > 0.$$

- 9. Find a non-constant function  $f \in \mathcal{A}(B(0,1))$  such that f has infinitely many zeros in B(0,1).
- 10. Suppose  $\alpha \neq 0$  is a root of a polynomial *p* of degree *n* with rational coefficients. Prove that  $1/\alpha$  is a root of a polynomial of degree at most *n* with rational coefficients.