

**Complex Variables**  
**Preliminary Exam**  
August 2013

**Directions:** Do all of the following eight problems. **Show all your work and justify your answers.** Each problem is worth 10 points.

**Notation:**  $\mathbb{C}$  — the complex plane;  $\mathbb{Z}$  — the set of integers;  $\mathbb{D} := \{z : |z| < 1\}$  — the unit disk;  $\Re(z)$  and  $\Im(z)$  denote the real part of  $z$  and the imaginary part of  $z$ , respectively.

1. Let  $z_1, z_2, \dots, z_n$  be  $n \geq 3$  distinct points in the complex plane which do not lie on a straight line. Prove that if

$$z_1 + z_2 + \dots + z_n = 0$$

then no closed half-plane, the boundary of which is a straight line passing through the origin, can contain all of the points  $z_1, z_2, \dots, z_n$ .

2. (a) Let  $f(z)$  be analytic in  $\mathbb{D}$  such that  $f(1/n) = f(-1/n)$  for all integers  $n \geq 2$ . Prove that  $f(z)$  is an even function.

(b) Let  $g(z)$  be analytic in the domain  $D := \{z : 0 < |z| < 1\}$ . Suppose that  $g(z) = g(2z)$  for all  $z$  such that  $|z| < 1/2$ . Prove that  $g(z)$  is constant.

3. Let

$$f(z) = \frac{\pi}{\sinh(\pi z)} + e^{1/z^2} + \frac{2z}{1+z^2}.$$

Locate and classify all the singularities of  $f(z)$  (including any singularity at  $z = \infty$ ) as isolated or non-isolated. Further, classify the isolated singularities by type (removable, pole, essential). Calculate the residues of  $f(z)$  at its poles.

4. Let  $f(z)$  be analytic in the strip  $S := \{z : |\Re(z)| < \pi/4\}$ . Suppose that  $f(0) = 0$  and  $|f(z)| < 1$  for all  $z \in S$ . Prove that  $|f(z)| \leq |\tan z|$  for all  $z \in S$ .

5. Let

$$f_n(z) = 1 + \frac{1}{z} + \frac{1}{2!z^2} + \dots + \frac{1}{n!z^n}.$$

Prove that for every  $\rho > 0$  there is positive integer  $N$  such that for all  $n \geq N$  all zeros of  $f_n(z)$  belong to the disk  $\{z : |z| < \rho\}$ .

6. Use a Residue Theorem to show that

$$\int_{-1}^1 \frac{\sqrt{1-x^2}}{1+x^2} dx = \pi(\sqrt{2}-1).$$

7. Find the number of solutions of the equation

$$z^4 + 2z^3 + 3z^2 + z + 2 = 0$$

in the right half-plane and in the first quadrant.

8. (a) State the Weierstrass Product Theorem.

(b) Give an example of a function  $f(z)$  analytic on  $\mathbb{C}$ , which has zeros at the points  $z_n = \frac{2n-1}{2}$ ,  $n \in \mathbb{Z}$  and no other zeros.