



OCCASIONAL PAPERS

Museum of Texas Tech University

Number 382

19 April 2022

EVIDENCE FROM OWL PELLETS INDICATE THAT PRAIRIE VOLES (*MICROTUS OCHROGASTER*) ARE BROADLY DISTRIBUTED IN THE TEXAS PANHANDLE

EMILY R. TURPEN, SARAH C. VRLA, RAYMOND S. MATLACK, AND ROBERT D. BRADLEY

ABSTRACT

To investigate the distribution of *Microtus ochrogaster* (Prairie Voles) in the Texas Panhandle, owl pellets were collected from 123 owl roosts in 27 counties. Remains of Prairie Voles were found at 76 roosts in 20 counties. Fifty-four of the roost sites (located in Briscoe, Castro, Deaf Smith, Floyd, Gray, Hale, Hartley, Hockley, Hutchinson, Lamb, Moore, Parmer, Randall, and Swisher counties) represent new county records for *M. ochrogaster* in Texas. Further, these data indicate that individuals and populations of *M. ochrogaster* in this region are more common than previously documented.

Key words: county records, *Microtus ochrogaster*, prairie vole, small mammals, Texas Panhandle

INTRODUCTION

Microtus ochrogaster (Prairie Vole) is a medium-size mouse that is widely distributed throughout the grasslands of the central United States (Stalling 1990). In the Oklahoma Panhandle and southwestern Kansas, *M. ochrogaster* occurs along riparian corridors, drainage routes, and related floodplains (Reed and Choate 1988). In western Oklahoma, skeletal remains of *M. ochrogaster* have been recovered from owl pellets collected in Ellis, Harmon, and Jackson counties (which border the Texas Panhandle), as well as from 11 other counties in western Oklahoma, resulting in an expansion of the known range in that state (McDonald et al. 2006).

Regarding the Texas Panhandle, Manning and Jones (1988) and Choate and Killebrew (1991) docu-

mented *M. ochrogaster* in Lipscomb and Hansford counties, respectively. More than 10 years later, two specimens of *M. ochrogaster* were captured in Carson County and skeletal remains were identified in owl pellets from Dallam and Sherman counties in the Texas Panhandle (McCaffery et al. 2003). Poole and Matlack (2007) reported the presence of *M. ochrogaster* in Armstrong, Carson, and Ochiltree counties based on skeletal remains found in owl pellets. Most recently, Roberts et al. (2015) reported a record of the Prairie Vole from Lubbock County, extending the range of the species approximately 150 km to the south. Based on these published reports, Schmidly and Bradley (2016) listed the documented range of the Prairie Vole to include Armstrong, Hansford, Lipscomb, and Lubbock counties in the Texas Panhandle.

Studies by Smith (1992) and Dalquest et al. (1990) in New Mexico, Oklahoma, and Texas suggested that livetrapping of *M. ochrogaster* in this portion of the prairie vole's range was difficult and they reported low rates of capture over many trap nights. Poole and Matlack (2007) suggested that populations of *M. ochrogaster* in the Panhandle region are ephemeral, also making sampling difficult. However, it is well known that owl pellets often offer insight into species of small mammals that are difficult to trap and owl pellets thereby offer a short-term archive of the small mammals in an area. For example, Wright et al. (2016) documented new county records of two genera

of shrews (*Notiosorex* and *Cryptotis*) from the Texas Panhandle that were unknown prior to the examination of a series of owl pellets.

Herein, owl pellets were collected and skeletal remains analyzed from multiple locations across the Texas Panhandle. Owl pellets from these locations often were found in sheltered/protected locations (e.g., barns or silos); consequently, they represent an accumulation of material over a number of years. The species of owls observed roosting across these locations were the Barn Owl (*Tyto alba*) and the Great Horned Owl (*Bubo virginianus*).

METHODS AND MATERIALS

Owl pellets were collected from Armstrong, Bailey, Briscoe, Carson, Castro, Cottle, Dallam, Deaf Smith, Donley, Floyd, Gray, Hale, Hall, Hansford, Hartley, Hockley, Hutchinson, Lamb, Lipscomb, Moore, Ochiltree, Oldham, Parmer, Randall, Sherman, Swisher, and Wheeler counties in the Panhandle of Texas. Cottle County is a peripheral county of the Texas Panhandle and Rolling Plains regions. Pellets were dissolved in an 8% NaOH (sodium hydroxide) solution for 20–24 hours to remove hair and release skeletal material from the tightly bound pellet. Skeletal

material was rinsed with water in a sieve and air dried for 24–48 hours. Individuals of *M. ochrogaster* were identified to species by cranial and dental characteristics (Elbroch 2006). Manning and Jones (1988), Choate and Killebrew (1991), McCaffrey et al. (2003), Poole and Matlack (2007), and Schmidly and Bradley (2016) were referenced to determine distribution within the known range of *M. ochrogaster* and new county records. All specimens obtained in the study were deposited in the Natural Science Research Laboratory, Museum of Texas Tech University (see Appendix).

RESULTS

Approximately 3,911 complete and numerous partial owl pellets and scattered skeletal remains were collected near 123 owl roosts in 27 counties in the Texas Panhandle. Remains of *M. ochrogaster* (skulls or jaws; 648 specimens) were identified at 76 roosts

in 20 counties. Of these, Briscoe, Castro, Deaf Smith, Floyd, Gray, Hale, Hartley, Hockley, Hutchinson, Lamb, Moore, Parmer, Randall, and Swisher counties represent new county records for the Prairie Vole in Texas (Fig. 1 and Appendix).




Figure 1. Location of owl roosts surveyed in the Texas Panhandle. The numbers placed within a county indicate the number of owl roosts wherein remains of *Microtus ochrogaster* were discovered. Refer to the Appendix for additional details and localities within counties. An asterisk (*) denotes county records for the species.

DISCUSSION

Prior to this study, *M. ochrogaster* was known to occur only in a few counties in the northern and north-central portions of the Texas Panhandle, with a single record from south of the Panhandle in Lubbock County (Manning and Jones 1988; Choate and Killebrew 1991; McCaffrey et al. 2003; Poole and Matlack 2007; Roberts et al. 2015; Schmidly and Bradley 2016). Remains of *M. ochrogaster* obtained from owl pellets indicate that the species is broadly distributed and perhaps common in the western and central portions of Texas Panhandle. The new information discovered in this study supports the statement of Schmidly and Bradley (2016) that *M. ochrogaster* may be expanding its range in Texas. This theory is supported by previous

studies conducted in 1987 through 2015 that reported few accounts of *M. ochrogaster* throughout the Texas Panhandle via trapping and the examination of owl pellets (Jones et al. 1987, 1988; Pesaturo 1989; Manning and Jones 1990; Goetze et al. 1991; Yancey et al. 1998; Lockwood and Jones 2000; Jones et al. 2015). Additional sampling of owl pellets in eastern New Mexico and additional portions of Texas, including the eastern Panhandle and areas to the south and east, is needed to better understand the distribution of the prairie vole in this portion of their range. In this regard, collection and examination of owl pellets offers an efficient method for surveying small mammal communities and providing distributional data for lesser-known species.

ACKNOWLEDGMENTS

We thank numerous landowners for access to roost sites. H. J. Garner provided advice and expertise in creating the catalog of specimens collected during

this study. West Texas A&M University President's Undergraduate Research Program grant to E.R.T. provided funding for this study.

LITERATURE CITED

- Choate, L. L., and F. C. Killebrew. 1991. Distributional records of the California myotis and the prairie vole in the Texas Panhandle. *Texas Journal of Science* 43:214–215.
- Dalquest, W. W., F. B. Stangl, Jr., and J. K. Jones, Jr. 1990. Mammalian zoogeography of a Rocky Mountain-Great Plains interface in New Mexico, Oklahoma, and Texas. *Special Publications, Museum of Texas Tech University* 34:1–78.
- Elbroch, M. 2006. *Animal skulls: a guide to North American species*, 1st ed. Stackpole Books, Mechanicsburg, Pennsylvania.
- Goetze, J. R., J. K. Jones, Jr., and R. W. Manning. 1991. Mammal remains from owl pellets from two localities in Swisher County, Texas. *Texas Journal of Science* 43:221–222.
- Jones, Jr., J. K., R. W. Manning, R. R. Hollander, and C. Jones. 1987. Annotated checklist of recent mammals of northwestern Texas. *Occasional Papers, Museum of Texas Tech University* 111:1–14.
- Jones, Jr., J. K., R. W. Manning, C. Jones, and R. R. Hollander. 1988. Mammals of the northern Texas Panhandle. *Occasional Papers, Museum of Texas Tech University* 126:1–54.
- Jones, C., F. D. Yancey, II, and R. W. Manning. 2015. The mammals of Caprock Canyon State Park, Texas. *Occasional Papers, Museum of Texas Tech University* 335:1–26.
- Lockwood, M. W., and C. Jones. 2000. Vertebrate remains found in owl pellets from Crosby County, Texas. *Texas Journal of Science* 52:169–173.
- Manning, R. W., and J. K. Jones, Jr. 1988. A specimen of the prairie vole, *Microtus ochrogaster*, from the northern Texas Panhandle. *Texas Journal of Science* 40:463–464.
- Manning, R. W., and J. K. Jones, Jr. 1990. Remains of small mammals recovered from barn owl pellets from Crosby County, Texas. *Texas Journal of Science* 42:311–312.
- McCaffrey, R. E., M. C. Wallace, J. F. Kamler, and J. D. Ray. 2003. Noteworthy distributional records of the prairie vole in the Texas and Oklahoma Panhandles. *The Southwestern Naturalist* 48:717–719.
- McDonald, B. K., P. W. Wilson, and W. Caire. 2006. New county records of Oklahoma mammals based on remains identified in owl pellets. *Proceedings of the Oklahoma Academy of Science* 86:47–52.
- Pesaturo, R. J., R. W. Manning, and J. K. Jones, Jr. 1989. Small mammals captured by barn owls in Lamb County, Texas. *Texas Journal of Science* 41:433–434.
- Poole, M. W., and R. S. Matlack. 2007. Prairie vole and other small mammals from the Texas Panhandle. *The Southwestern Naturalist* 52:442–445.
- Reed, M. P., and J. R. Choate. 1988. Noteworthy southwestern records of the prairie vole. *The Southwestern Naturalist* 33:495–496.
- Roberts, E. K., et al. 2015. A record of *Microtus ochrogaster* from the Llano Estacado and other distributional records of small mammals from Texas. *Occasional Papers, Museum of Texas Tech University* 329:1–7.
- Schmidly, D. J., and R. D. Bradley. 2016. *The mammals of Texas*, 7th ed. University of Texas Press, Austin.
- Smith, K. S. 1992. The prairie vole, *Microtus ochrogaster*, in Caddo County, Oklahoma. *Texas Journal of Science* 44:116–117.
- Stalling, D. T. 1990. *Microtus ochrogaster*. *Mammalian Species* 355:1–9.
- Wright, E. A., E. K. Brookover, B. A. Gross, J. Q. Francis, and R. D. Bradley. 2016. Noteworthy records of shrews from the Panhandle of Texas. *Occasional Papers, Museum of Texas Tech University* 342:1–3.
- Yancey, F. D., R. W. Manning, J. R. Goetze, and C. Jones. 1998. The mammals of Lake Meredith National Recreational Area and adjacent areas, Hutchinson, Moore, and Potter Counties, Texas. *Occasional Papers, Museum of Texas Tech University* 174:1–20.

Addresses of authors:

EMILY R. TURPEN

*PO Box 1834
Hereford, TX 79045
emturpen1546@gmail.com*

RAYMOND S. MATLACK

*West Texas A&M University
Box 60808
Canyon, TX 79016
rmatlack@wtamu.edu*

SARAH C. VRLA

*Department of Biological Sciences
Texas Tech University
Lubbock, TX 79409
sarah.vrla@ttu.edu*

ROBERT D. BRADLEY

*Department of Biological Sciences and
Natural Science Research Laboratory
Museum of Texas Tech University
Lubbock, TX 79409
robert.bradley@ttu.edu*

Associate Editor for this manuscript was Richard D. Stevens

APPENDIX

Microtus ochrogaster specimens collected in this study are reported below. Counties are listed alphabetically and are followed by number of specimens per county (in parentheses). This is followed by the number of specimens per locality, latitude/longitude data, and Museum of Texas Tech University catalog number (in parentheses).

Specimens examined.—Armstrong County (15): 8, 35°06'14.0"N, 101°34'04.0"W (TTU-M 150618–150625); 2, 35°06'20.0"N, 101°28'51.0"W (TTU-M 150626–150627); 3, 34°59'40.6"N, 101°20'21.5"W (TTU-M 150628–150630); 2, 35°02'25.2"N, 101°22'06.1"W (TTU-M 150631–150632).

Briscoe County (32): 3, 34°42'22.8"N, 101°26'29.7"W (TTU-M 150633–150635); 4, 34°28'46.0"N, 101°23'42.5"W (TTU-M 150636–150639); 18, 34°24'17.2"N, 101°26'07.7"W (TTU-M 150640–150657); 7, 34°27'53.8"N, 101°13'00.9"W (TTU-M 150658–150664).

Carson County (58): 9, 35°12'07.6"N, 101°15'30.8"W (TTU-M 150665–150673); 12, 35°15'45.7"N, 101°14'59.6"W (TTU-M 150674–150685); 22, 35°26'31.8"N, 101°16'48.4"W (TTU-M 150686–150707); 4, 35°19'27.8"N, 101°17'34.2"W (TTU-M 150708–150711); 11, 35°20'29.1"N, 101°14'24.4"W (TTU-M 150712–150722).

Castro County (76): 2, 34°20'50.0"N, 102°18'05.0"W (TTU-M 150723–150724); 5, 34°22'31.0"N, 102°18'14.0"W (TTU-M 150725–150729); 29, 34°42'11.5"N, 102°00'51.9"W (TTU-M 150730–150758); 3, 34°41'16.5"N, 102°09'32.1"W (TTU-M 150759–150761); 36, 34°41'19.2"N, 102°12'11.1"W (TTU-M 150762–150797); 1, 34°33'25.9"N, 102°14'29.4"W (TTU-M 150798).

Dallam County (2): 2, 36°09'14.7"N, 102°26'08.7"W (TTU-M 150799–150800).

Deaf Smith County (20): 9, 34°54'34.0"N, 102°19'53.0"W (TTU-M 150801–150809); 7, 34°45'23.4"N, 102°31'06.3"W (TTU-M 150810–150816); 2, 34°47'02.5"N, 102°30'50.1"W (TTU-M 150817–150818); 2, 35°06'34.3"N, 102°12'08.6"W (TTU-M 150819–150820).

Floyd County (28): 4, 34°12'49.9"N, 101°26'01.1"W (TTU-M 150821–150824); 11, 34°07'29.7"N, 101°25'51.4"W (TTU-M 150825–150835); 2, 33°58'33.6"N, 101°28'29.2"W (TTU-M 150836–150837); 4, 33°52'49.8"N, 101°30'51.1"W (TTU-M 150838–150841); 7, 34°04'55.6"N, 101°17'49.2"W (TTU-M 150842–150848).

Gray County (9): 2, 35°23'59.6"N, 100°58'07.7"W (TTU-M 150849–150850); 7, 35°19'44.2"N, 101°03'11.1"W (TTU-M 150851–150857).

Hale County (54): 16, 34°14'59.2"N, 101°39'21.3"W (TTU-M 150858–150873); 11, 33°56'30.7"N, 101°41'42.7"W (TTU-M 150874–150884); 5, 33°52'01.5"N, 101°44'10.2"W (TTU-M 150885–150889); 8, 33°52'37.8"N, 101°50'47.1"W (TTU-M 150890–150897); 2, 33°55'54.2"N, 102°00'41.1"W (TTU-M 150898–150899); 8, 34°04'01.1"N, 101°55'24.9"W (TTU-M 150900–150907); 4, 34°12'00.2"N, 102°05'03.7"W (TTU-M 150908–150911).

Hansford County (55): 1, 36°08'21.9"N, 101°12'43.7"W (TTU-M 150912); 3, 36°11'57.8"N, 101°07'41.2"W (TTU-M 150913–150915); 2, 36°19'27.9"N, 101°16'27.2"W (TTU-M 150916–150917); 11, 36°20'36.1"N, 101°31'03.5"W (TTU-M 150918–150928); 38, 36°17'02.2"N, 101°31'53.9"W (TTU-M 150929–150966).

Hartley County (2): 2, 36°01'40.1"N, 102°17'37.8"W (TTU-M 150967–150968).

Hockley County (8): 6, 33°47'13.0"N, 102°13'39.0"W (TTU-M 150969–150974); 2, 33°46'13.0"N, 102°13'37.0"W (TTU-M 150975–150976).

Hutchinson County (56): 31, 35°56'56.5"N, 101°27'32.0"W (TTU-M 150977–151007); 13, 36°01'21.1"N, 101°19'09.1"W (TTU-M 151008–151020); 12, 35°59'17.2"N, 101°13'08.2"W (TTU-M 151021–151032).

Lamb County (7): 4, 33°55'37.2"N, 102°12'23.7"W (TTU-M 151033–151036); 3, 34°14'03.5"N, 102°31'35.8"W (TTU-M 151037–151039).

Moore County (34): 19, 35°52'14.0"N, 101°57'02.0"W (TTU-M 151040–151058); 15, 35°53'32.9"N, 101°39'07.1"W (TTU-M 151059–151073).

Ochiltree County (69): 3, 36°11'16.6"N, 100°49'12.4"W (TTU-M 151074–151076); 45, 36°16'57.5"N, 100°43'43.9"W (TTU-M 151077–151121); 20, 36°23'02.6"N, 100°43'25.7"W (TTU-M 151122–151141); 1, 36°11'39.8"N, 100°53'31.3"W (TTU-M 151142).

Parmer County (10): 1, 34°43'08.0"N, 102°32'30.0"W (TTU-M 151143); 3, 34°38'28.6"N, 102°48'15.6"W (TTU-M 151144–151146); 6, 34°31'32.7"N, 102°43'56.3"W (TTU-M 151147–151152).

Randall County (52): 2, 35°06'47.0"N, 101°44'02.0"W (TTU-M 151153–151154); 16, 34°54'31.7"N, 101°58'24.3"W (TTU-M 151155–151170); 9, 34°51'12.1"N, 101°53'06.0"W (TTU-M 151171–151179); 14, 34°50'05.6"N, 101°53'22.92"W (TTU-M 151180–151193); 4, 34°47'08.8"N, 102°00'35.2"W (TTU-M 151194–151197); 7, 34°47'32.0"N, 101°48'07.5"W (TTU-M 151198–151204).

Sherman County (12): 2, 36°10'56.4"N, 101°45'06.8"W (TTU-M 151205–151206); 3, 36°19'38.2"N, 101°44'33.2"W (TTU-M 151207–151209); 7, 36°19'39.8"N, 101°48'07.4"W (TTU-M 151210–151216).


Swisher County (49): 3, 34°31'41.3"N, 101°55'57.6"W (TTU-M 151217–151219); 4, 34°37'44.8"N, 101°45'28.2"W (TTU-M 151220–151223); 12, 34°38'45.3"N, 101°37'00.6"W (TTU-M 151224–151235); 2, 34°32'05.7"N, 101°35'34.9"W (TTU-M 151236–151237); 4, 34°27'46.8"N, 101°33'50.0"W (TTU-M 151238–151241); 12, 34°25'29.4"N, 101°34'32.6"W (TTU-M 151242–151253); 12, 34°22'34.1"N, 101°34'24.7"W (TTU-M 151254–151265).

PUBLICATIONS OF THE MUSEUM OF TEXAS TECH UNIVERSITY

This publication is available free of charge in PDF format from the website of the Natural Science Research Laboratory, Museum of Texas Tech University (www.depts.ttu.edu/nsrl). The authors and the Museum of Texas Tech University hereby grant permission to interested parties to download or print this publication for personal or educational (not for profit) use. Re-publication of any part of this paper in other works is not permitted without prior written permission of the Museum of Texas Tech University.

Institutional subscriptions to Occasional Papers are available through the Museum of Texas Tech University, attn: NSRL Publications Secretary, Box 43191, Lubbock, TX 79409-3191. Individuals may also purchase separate numbers of the Occasional Papers directly from the Museum of Texas Tech University.

Series Editor: Robert D. Bradley
Production Editor: Lisa Bradley
Copyright: Museum of Texas Tech University



ISSN 0149-175X

Museum of Texas Tech University, Lubbock, TX 79409-3191