

OCCASIONAL PAPERS

THE MUSEUM

TEXAS TECH UNIVERSITY

NUMBER 39

16 APRIL 1976

A NEW SPECIES OF *CHIRODERMA* FROM GUADELOUPE, WEST INDIES (CHIROPTERA: PHYLLOSTOMATIDE)

ROBERT J. BAKER AND HUGH H. GENOWAYS

During the course of a study of the bat faunas of the Caribbean islands we obtained a specimen of *Chiroderma* from the island of Guadeloupe in the Lesser Antilles. The nearest known populations of this genus occur on Tobago and Trinidad (*C. villosum* and *C. trinitatum*), approximately 550 kilometers to the south. This specimen represents a distinct new species that appears to be most closely related to *C. doriae* and *C. villosum*. This new species is named and described below.

***Chiroderma improvisum*, new species**

Holotype.—Adult male, skin and skull, number 19900 of The Museum, Texas Tech University (TTU); from Guadeloupe: Basse-Terre; 2 km. S, 2 km. E Baie-Mahault; obtained on 29 July 1974 by Genoways and Baker; original number John C. Patton 552; karyotype number TK 8285. Live cell lines are frozen in liquid nitrogen at The Museum, Texas Tech University.

Geographic distribution.—Presently known only from the type locality.

Description.—Size largest for genus, both externally and cranially (Table 1 and Figs. 1 through 4). Dorsal coloration grayish brown with a distinct white line down the center of the posterior half of the back; ventral coloration gray, tips of the hairs with a white band producing a "frosted" effect; indistinct white lines above and below each eye. Karyotypic features include a diploid number of 26 and a fundamental number of 48 (Fig. 5). The autosomes consist of eleven pairs

TABLE 1.—Measurements of five species of *Chiroderma*. See text for additional comparison of *C. improvisum* to *C. doriae*. All specimens are adults. Definition of measurements is given in Baker (1972). All measurements are in millimeters.

Museum number	Sex	Locality	Total length	Hind foot	Ear	Tragus	Forearm	Greatest length of skull	Condylbasal length	Zygomatic breadth	Mastoid breadth	Interorbital breadth	Breadth across upper canines	Maxillary tooth row	Mandibular length
19900	♂	Guadeloupe	87	15	21	7	57.5	29.9	27.7	18.9	14.3	6.5	7.2	10.7	20.3
<i>Chiroderma improvisum</i>															
5276	♂	Trinidad	73	11	17	6	47.1	25.3	22.2	15.7	11.9	5.8	6.0	8.5	16.0
9016	♂	Trinidad	76	11	18	5	47.3	25.6	22.9	15.1	12.1	5.8	6.0	8.5	16.3
9017	♀	Trinidad	66	11	15	4	46.6	26.0	22.5	15.8	12.2	5.9	6.1	8.9	16.3
5354	♀	Trinidad	75	12	18	7	48.9	26.2	23.5	17.0	12.9	6.2	6.2	9.2	16.9
5361	♀	Trinidad	64	11	17	6	46.8	24.9	22.7	15.3	11.9	5.6	6.0	8.8	15.6
<i>Chiroderma villosum</i>															
<i>Chiroderma salvini</i>															
9998	♂	Veracruz	65	12	19	3	47.0	25.2	22.3	15.5	12.0	5.7	6.0	8.6	15.9
9999	♂	Veracruz	62	11	15	5	46.2	24.4	22.0	14.0	11.2	5.5	5.7	8.5	15.5
12087	♂	Honduras	72	14	18	7	46.3	26.9	24.2	16.4	13.1	6.2	6.2	9.6	17.5
12800	♂	Honduras	76	13	19	8	47.5	26.5	24.1	16.1	12.5	6.2	6.0	9.3	17.0
6123	♂	Colima	72	11	17	7	43.7	24.4	21.8	15.0	11.8	5.8	5.2	8.6	15.3

TABLE 1.—Continued.

<i>Chiroderma triniatum</i>															
5223	♀	Trinidad	69	12	18	7	41.8	22.8	19.8	13.4	11.0	5.4	5.1	7.8	14.2
5382	♀	Trinidad	55	11	16	6	39.4	22.4	19.6	13.7	11.2	5.4	4.9	7.4	14.0
5487	♂	Trinidad	56	10	15	6	38.9	22.5	19.8	13.5	11.1	5.5	4.8	7.6	13.4
8989	♂	Trinidad	56	11	16	6	40.6	22.2	19.8	13.6	11.0	5.2	4.9	7.5	14.1
9014	♂	Trinidad	62	10	16	6	39.4	22.1	19.1	13.4	10.8	5.5	5.1	7.4	13.8
<i>Chiroderma doriei</i>															
Males*															
Brazil															
Average			74.8		20.2	7.3	52.0	28.0	26.1	17.6	13.8	6.3	6.4	10.2	19.8
Minimum			69.0		19.0	7.0	49.4	27.3	25.5	17.2	13.6	5.9	6.2	9.9	19.3
Maximum			78.5		21.5	7.5	53.5	28.7	26.8	18.5	14.3	6.7	6.6	10.4	20.2
Females**															
Brazil															
Average			75.5		20.4	7.3	53.0	28.2	26.3	17.8	14.0	6.3	6.5	10.3	19.8
Minimum			70.0		19.0	7.0	51.0	27.5	25.6	16.9	13.6	6.1	6.1	10.1	19.4
Maximum			80.0		21.5	7.5	55.5	28.7	26.7	18.4	14.3	6.6	6.8	10.5	20.3

*Sample size 15 (Taddei, 1973).

**Sample size 21 for external measurements and 15 for cranial measurements (Taddei, 1973).



1




FIG. 1.—Photograph of the face and venter of the holotype of *Chiroderma improvisum* in life.

FIG. 2.—Anterior view of the upper incisors and canines of four species of *Chiroderma*: A, *C. salvini*; B, *C. trinitatum*; C, *C. villosum*; and D, *C. improvisum*.




FIG. 3.—Dorsal views of the skulls of four species of *Chiroderma*: A, *C. salvini*; B, *C. trinitatum*; C, *C. villosum*; and D, *C. improvisum*. Bar represents 5 millimeters.

FIG. 4.—Dorsal, ventral, and lateral views of the skull of the holotype of *Chiroderma improvisum*. Bar represents 5 millimeters.




FIG. 5.—A, karyotype of the holotype of *Chiroderma improvisum*; B, karyotype of a male *Chiroderma trinitatum* from Trinidad.

of biarmed elements and one pair of acrocentric elements (the smallest pair of autosomes). Of the 11 biarmed pairs of autosomes, two pairs have a subtelocentric placement of the centromere whereas the remainder have either a submetacentric or metacentric placement of the centromere. The X-chromosome is subtelocentric and the Y-chromosome is a small element with a minute second arm.

Measurements.—Measurements for the holotype of *C. improvisum* and the four other species of *Chiroderma* are given in Table 1. The holotype of *C. doriae* (the species nearest *C. improvisum* in size) was measured by Dr. Dilford C. Carter during a visit to the British Museum (Natural History) in 1966, and he has kindly allowed us to use his measurements. The skull of the holotype of *doriae* is broken, and the forearms were damaged in preparation so that only a few measurements could be taken. For consistency, we asked Dr. Carter to take the same measurements on the holotype of *C. improvisum*.

His measurements for the holotypes of *C. improvisum* and *C. doriae*, respectively, follow: metacarpal III, 57.6, 52.8; phalanx 1, digit III, 22.8, 20.5; phalanx 2, digit III, 31.0, 28.5; metacarpal IV, 54.9, 51.7; phalanx 1, digit IV, 19.2, 17.1; metacarpal V, 46.8,

53.4; phalanx 1, digit V, 24.8, 12.9; length of tibia, 19.8, 18.0; post-orbital constriction, 6.5, 6.2; length of mandible, 21.2, 18.9; length of mandibular toothrow, 12.1, 11.5; length of maxillary toothrow, 11.0, 10.6; breadth across upper molars, 13.6, 13.2; breadth across upper canines, 7.2, 6.6; length of nose leaf, 5.9, 6.0; length of calcar, 7.2, 6.2.

Additional measurements are given for species of *Chiroderma* in Goodwin (1946:321-322), Goodwin and Greenhall (1961:257-259), Handley (1960, 1965), Husson (1962:166-169), and Villa-R. (1966:288-293).

Comparison.—*Chiroderma improvisum* is the largest species of the genus. This is reflected in the 13 measurements in Table 1, where the only overlap between *C. improvisum* and the other three common species is in length of tragus. Measurements for the holotype of *C. improvisum* are larger than any recorded for *C. doriae* (Taddei, 1973) in total length, forearm, greatest length of skull, condylobasal length, zygomatic breadth, breadth across upper canines, and length of maxillary toothrow (Table 1). Other measurements of *C. improvisum* are large relative to the range reported for *C. doriae* (Taddei, 1973). *Chiroderma improvisum* has more massive canines (Fig. 2) than any of the three more common species. Upper incisor one is quite broad at the cingulum, and the outside lateral cingulum is better developed than in *C. villosum*, *C. salvini*, or *C. trinitatum*. The holotype of *C. improvisum* has a distinct white line down the back. Such a dorsal marking is recorded as absent from the holotype of *C. doriae*. However, the specimens of Taddei (1973, figs. 2-6, 33, and personal communication) show a broad dorsal stripe and white stripes above and below the eye. These stripes in Taddei's specimens are more distinct than those of the holotype of *C. improvisum*.

Chromosomally, the karyotype of *C. improvisum* (Fig. 5) is distinguished from that of *C. villosum*, *C. trinitatum* (Baker, 1970), and *C. salvini* (Baker, 1973) by the presence of a pair of small, nearly acrocentric autosomes and only two pairs of subtelocentric autosomes. In *C. villosum*, *C. salvini*, and *C. trinitatum*, there are three pairs of autosomes with a subtelocentric placement of the centromere, and the smallest pair of autosomes is submetacentric or metacentric in nature.

Remarks.—*Chiroderma improvisum* is obviously specifically distinct from *C. villosum*, *C. salvini*, and *C. trinitatum*. We have not had the opportunity to examine a specimen of *C. doriae*, but from Table 1 it is clear that the holotype of *C. improvisum* is considerably larger than any specimen of *C. doriae* thus far reported. It is possible that *C. improvisum* and *C. doriae* are relictual populations of a once

widespread species of *Chiroderma* and that their large size and similar descriptions of canines and incisors (Thomas, 1891) reflect a true relationship. Of the species on Guadeloupe, only *Artibeus jamaicensis* is found in the area of southeastern Brazil where *C. doriae* has been reported, and it seems unlikely that populations of only a single species would be found on Guadeloupe and in southeastern Brazil. Another possibility is that these two taxa represent the product of convergent evolution. Some taxa of bats (*Eptesicus guadeloupensis* and *Sturnira thomasi*) on Guadeloupe are larger than their nearest evolutionary relatives. If convergent evolution has accounted for this similarity between *C. doriae* and *C. improvisum*, then the nearest evolutionary relative of *C. improvisum* is probably *C. villosum*. Karyotypic data from *C. doriae* would certainly be valuable, for if *C. doriae* had the *C. improvisum* karyotype, a common ancestor for these two forms would be highly probable.

Etymology.—The specific name *improvisum* is from Latin, meaning unforeseen or unexpected. We did not expect to find *Chiroderma* on Guadeloupe because the nearest known representative from the Caribbean occurs on Trinidad and Tobago, 550 kilometers to the south.

Acknowledgments.—We thank Dr. Dillard C. Carter for allowing us to use his measurements of *C. doriae* and for measuring the holotype of *C. improvisum*. We thank John W. Bickham and John C. Patton for assistance in the field. Stephen L. Williams and Laura Kyle assisted in preparation of figures. Drs. William B. Davis and Karl F. Koopman critically evaluated the manuscript. Supported by National Science Foundation grant GB-41105 and the Institute of Museum Research, Texas Tech University.

LITERATURE CITED

- BAKER, R. J. 1970. Karyotypic trends in bats. Pp. 65-96, in *Biology of bats* (W. A. Wimsatt, ed.), Academic Press, New York, 1:xii + 1-406.
- . 1973. Comparative cytogenetics of New World leaf-nosed bats (Phyllostomatidae). *Periodicum biologorum*, 75:37-45.
- GOODWIN, G. G. 1946. Mammals of Costa Rica. *Bull. Amer. Mus. Nat. Hist.*, 87:275-473.
- GOODWIN, G. G., AND A. M. GREENHALL. 1961. A review of the bats of Trinidad and Tobago. *Bull. Amer. Mus. Nat. Hist.*, 122:187-302.
- HANDLEY, C. O., JR. 1960. Descriptions of new bats from Panama. *Proc. U.S. Nat. Mus.*, 112:459-479.
- . 1965. Descriptions of new bats (*Chiroderma* and *Artibeus*) from Mexico. *An. Inst. Biol., Univ. Nac. México*, 36:297-301.
- HUSSON, A. M. 1962. The bats of Suriname. *Zool. Verhandl.*, 58:1-282.

- TADDEI, V. A. 1973. Phyllostomidae da Região Norte-Occidental do Estado de São Paulo. Unpublished Ph.D. thesis, Fac. Filos. Ciênc. Letr. São José do Rio Preto, S. P. (Brazil).
- THOMAS, O. 1891. Note on *Chiroderma villosum*, Peters, with the description of a new species of the genus. *Ann. Mus. Civ. Stor. Nat. Genova*, 10:881-883.
- VILLA-R., B. 1966. Los murciélagos de México. *Inst. Biol., Univ. Nac. Auto. México*, xvi+491 pp.

Addresses of authors: ROBERT J. BAKER, *The Museum and Department of Biological Sciences*; HUGH H. GENOWAYS, *The Museum, Texas Tech University, Lubbock, 79409*. Received 23 April, accepted 6 June 1975.

PUBLICATIONS OF THE MUSEUM
TEXAS TECH UNIVERSITY

Two publications of The Museum of Texas Tech University are issued under the auspices of the Dean of The Graduate School and Director of Academic Publications, and in cooperation with the International Center for Arid and Semi-Arid Land Studies. Shorter research papers are published as Occasional Papers, whereas longer contributions appear as Special Publications. Both are numbered separately and published on an irregular basis.

Institutional libraries interested in exchanging publications may obtain the Occasional Papers and Special Publications by addressing the Exchange Librarian, Texas Tech University, Lubbock, Texas 79409. Individuals may purchase separate numbers of the Occasional Papers for 50 cents each through the Exchange Librarian. Remittance must be enclosed with request.