RESEARCH ARTICLE

Open Access

Mutation in a PHD-finger protein MS4 causes male sterility in soybean



Sandi Win Thu¹⁺, Krishan Mohan Rai¹⁺, Devinder Sandhu^{2*}, Alex Rajangam³, Vimal Kumar Balasubramanian¹, Reid G. Palmer^{4^} and Venugopal Mendu^{1*}

Abstract

Background: Male sterility has tremendous scientific and economic importance in hybrid seed production. Identification and characterization of a stable male sterility gene will be highly beneficial for making hybrid seed production economically feasible. In soybean, eleven male-sterile, female-fertile mutant lines (*ms1, ms2, ms3, ms4, ms5, ms6, ms7, ms8, ms9, msMOS*, and *msp*) have been identified and mapped onto various soybean chromosomes, however the causal genes responsible for male sterility are not isolated. The objective of this study was to identify and functionally characterize the gene responsible for the male sterility in the *ms4* mutant.

Results: The *ms4* locus was fine mapped to a 216 kb region, which contains 23 protein-coding genes including *Glyma.02G243200*, an ortholog of Arabidopsis *MALE MEIOCYTE DEATH 1 (MMD1)*, which is a Plant Homeodomain (PHD) protein involved in male fertility. Isolation and sequencing of *Glyma.02G243200* from the *ms4* mutant line showed a single base insertion in the 3rd exon causing a premature stop codon resulting in truncated protein production. Phylogenetic analysis showed presence of a homolog protein (MS4_homolog) encoded by the *Glyma.14G212300* gene. Both proteins were clustered within legume-specific clade of the phylogenetic tree and were likely the result of segmental duplication during the paleoploidization events in soybean. The comparative expression analysis of *Ms4* and *Ms4_homologs* across the soybean developmental and reproductive stages showed significantly higher expression of *Arabidopsis mmd1* mutant with the soybean *Ms4* gene produced normal stamens, successful tetrad formation, fertile pollens and viable seeds, whereas the *Ms4_homolog* was not able to restore male fertility.

Conclusions: Overall, this is the first report, where map based cloning approach was employed to isolate and characterize a gene responsible for the male-sterile phenotype in soybean. Characterization of male sterility genes may facilitate the establishment of a stable male sterility system, highly desired for the viability of hybrid seed production in soybean. Additionally, translational genomics and genome editing technologies can be utilized to generate new male-sterile lines in other plant species.

Keywords: Fertility, Mapping, mmd1, ms4, Plant homeodomain, Soybean, Sterility

Highlight

Fine mapping of soybean *ms4* resulted in identification of male sterile phenotype causal mutation. The MS4 is a PHD-finger protein which is functionally characterized using Arabidopsis *mmd1* mutant.

* Correspondence: devinder.sandhu@ars.usda.gov; venugopal.mendu@ttu.edu

¹Sandi Win Thu and Krishan Mohan Rai contributed equally to this work. ²US Salinity Laboratory (USDA-ARS), Riverside, CA 92507, USA ¹Fiber and Biopolymer Research Institute, Department of Plant and Soil Science, Texas Tech University, Lubbock, TX 79409, USA Full list of author information is available at the end of the article

Background

Soybean (*Glycine max*) is one of the most important food crops in the world known for its high seed protein and oil content [1]. In general, the crop performance and yield can be increased by generating hybrid plants which outcompete pure lines due to heterosis or hybrid vigor [2]. As soybean is a self-pollinating crop, producing large quantities of hybrid seed by manual emasculation and cross-pollination is difficult and economically nonviable. Hence, identification of a stable male sterility system similar to rice and maize is much needed in soybean [3]. Male sterility resulting from mutations in genes



© The Author(s). 2019 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated. involved in microsporogenesis and/or microgametogenesis has been described in various plant species [4]. Mutations in the genes involved in synapsis (proper chromosomal pairing and gamete formation) are known to be responsible for male-sterile female-sterile, malesterile female-fertile or male-fertile female-sterile phenotypes in plants [5]. In soybean, so far, a total of 11 malesterile, female-fertile (ms1, ms2, ms3, ms4, ms5, ms6, ms7, ms8, ms9, msMOS and msp) mutants have been identified and mapped on to different chromosomes [6-8]. Further, genetic inheritance and allelism tests, revealed that these 11 soybean male-sterile mutants are genetically independent of each other [7]. Cytological studies on these mutants suggested various mechanisms governing the male-sterile phenotype, which include cytokinesis failure during telophase II, failure of tetrad formation during microsporogenesis, microspore and pollen degeneration, abnormality/lack of vacuoles, and callose level **[4**, 7, 9–14]. Molecular low characterization of the male sterility genes will be the key to establish a sterility system for commercial application in hybrid seed production.

Among the male-sterile mutant lines, the *ms4* mutant was an outcome of a spontaneous mutation in cultivar "Rampage" that was identified at the Iowa State University in 1973 (Fig.1) [15]. Phenotyping of homozygous

ms4/ms4 plants showed that the anthers were slightly smaller in size and lighter in color compared to the control [15]. The ms4 mutant forms clumps of degenerated empty microspores due to the failure of cytokinesis after telophase II resulting in coenocytic microspores [10]. Genetic analysis of the ms4 mutant line revealed that the male-sterile phenotype is governed by a single recessive gene that mapped to a 694 kb region containing 88 predicted protein-coding genes on soybean chromosome 2 [6]. Fine mapping and/or causal gene identification of ms4 will help in designing molecular markers to screen male-sterile plants for hybrid seed production. The process of hybrid seed production requires planting pure male-sterile plants next to the donor soybean genotypes in field for cross pollination to occur. However, a malesterile line cannot be maintained in homozygous condition as it does not produce seed, hence, is maintained in heterozygous condition. For hybrid seed production, male-fertile plants (homozygous and heterozygous) that arise in the progeny of heterozygous male-sterile plants need to be eliminated to prevent self-pollination. Identification of the causal male-sterility gene can facilitate marker-based genotyping of the segregating male-sterile population to eliminate fertile plants during the seedling stage before flowering. In addition, the functional characterization of male-sterile genes will also help to

Image: A state of the state

elucidate the molecular mechanism responsible for the male sterility and extend the technology to other crops to develop male-sterile lines using genetic engineering.

In the present investigation, we have fine mapped the *Ms4* locus to a 216 kb region containing 23 predicted protein-coding genes, identified the candidate gene (*Gly-ma.02G243200*) based on the functional relevance and characterized it using Arabidopsis model system. The protein sequence analysis showed that *Ms4* (*Gly-ma.02G243200*) is an ortholog of Arabidopsis *MALE MEIOCYTE DEATH 1* (*MMD1*), a PHD-finger protein involved in male fertility. Further, functional characterization of *Glyma.02G243200* using Arabidopsis *mmd1* mutant restored the male-fertile phenotype in Arabidopsis, thereby confirming its role in male fertility.

Results

Gene discovery by fine mapping of the ms4 locus

We previously mapped the *ms4* locus on to chromosome 2 between the microsatellite markers Satt703 and BARC-SOYSSR_02_1539 [6]. This region physically spanned 694 kb and harbored 88 predicted protein-coding genes. To further fine map the *ms4* locus, 20 potential SSR (Simple Sequence Repeat) markers were used to test the polymorphism in the chromosomal region between Satt703 and BARCSOYSSR_02_1539 of the parental lines of the mapping population; T274H (*ms4/ms4*) as a female parent and Manchu (*Ms4/Ms4*) as a male parent. Of these, 8 markers (BARCSOYSSR_02_1509, BARCSOYSSR_02_1510, BARCSOYSSR_02_1513, BARCSOYSSR_02_1514, BARCSOYSSR_02_1515, BARCSOYSSR_02_1528, BARCSOYSSR_02_1529 and BARCSOYSSR_02_1530) showed polymorphism between the two parental lines. These 8

polymorphic markers were further used for fine mapping of the *ms4* locus using an F_2 population consisting of 118 plants. The *ms4* locus was mapped to 5.1 cM region between BARCSOYSSR_02_1515 and BARCSOYSSR_02_ 1528 (Fig. 2). The corresponding region on the soybean physical map was 216 kb with 23 predicted protein-coding genes (Fig. 2; Additional File 1: Table S1). A closer look at the functional annotation of 23 protein-coding genes showed a candidate gene of interest, *Glyma.02G243200*, which encodes a protein with 64% homology to Arabidopsis MALE MEIOCYTE DEATH 1 (MMD1) protein. AtMMD1 is involved in the chromosomal condensation process during the microsporogenesis, and mutations in *MMD1* leads to male sterility in Arabidopsis [16–18].

Identification of the *ms4* causal mutation by sequence comparison

The candidate gene of interest Glyma.02G243200 was successfully amplified from the genomic DNA of male-fertile (Ms4/Ms4) as well as homozygous male-sterile (ms4/ms4) soybean plants. Out of 9 different combinations of forward and reverse primers designed from 1,500 bp up- and down-stream sequences flanking the Glyma.02G243200 gene, only MS4_F2 and MS4_R1 primer pair was able to amplify the expected fragment of 4.554 kb comprising 3.1 kb Glyma.02G243200 gene (Fig. 3a; Additional File 1: Table S2A). The coding sequence comparison showed presence of three exons and two introns (Fig. 3a). The amplified fragment was sequenced with 12 internal primer pairs to cover the entire 3.1 kb region of Glyma.02G243200 (Additional File 1: Table S2B). The nucleotide sequence of Glyma.02G243200 from the male-sterile plant was compared with the male-fertile plants. The sequence



distances are shown in base pairs



comparison of the genomic DNA sequences showed an insertion of a single adenine (A) nucleotide in the 3rd exon in the male-sterile (*ms4/ms4*) line (Additional File 1: Figure S1). This single base insertion caused a frameshift mutation that resulted in a premature stop codon thereby forming a truncated protein of 430 amino acids (Fig. 3b; Additional File 1: Figure S2). Further, *Glyma.02G243200* transcripts from the wild type fertile and the *ms4* sterile mutants were also sequenced to validate the presence of the mutation and to confirm the exon-intron boundaries (data not shown).

Phylogenetic analysis showed highly conserved nature of the MS4 proteins

The bioinformatic analysis of the MS4 protein sequence showed that it is a 707 amino acid long protein (Fig. 3b) with a theoretical molecular weight of 79.65 KDa and theoretical pI of 5.97 [19]. Further, conserved domain analysis showed that MS4 protein is a plant homeodomain (PHD)finger protein as it contains a PHD_MMD1_like domain (CD15556) (Fig. 3b). The PHD domain is present at the C-terminus between 623 and 667 amino acid residues. The domain architecture shown by the soybean MS4 protein is consistent with its Arabidopsis counterpart PHD finger protein MALE MEIOCYTE DEATH 1 (AtMMD1). However, conserved domain analysis of the MS4 protein derived from male-sterile soybean line showed the absence of PHD domain in the truncated MS4 protein (Fig. 3b).

The phylogenetic analysis was performed with the MS4 protein and its homologs identified in other plant species through protein basic local alignment search tool (BLAST) of NCBI. The phylogenetic tree showed clustering of the proteins based on their taxonomical relationship (Fig. 4). The soybean MS4 (XP_003518369) showed phylogenetic

similarity with proteins derived from other legume species such as common bean (Phaseolus vulgaris; XP_007141934), mung bean (Vigna radiata; XP_014503023), adzuki bean (V. angularis; XP_017428580) and pigeon pea (Cajanus cajan; XP_020202556), etc. and all of them clustered into a Fabaceae specific cluster (Fig. 4). Surprisingly, we found another PHD domain containing protein (XP_003544360; MS4_homolog) in soybean with a high similarity with MS4 (Fig. 4). The MS4_homolog is a 708 aa long protein encoded by Glyma.14G212300 (Gene: 3178 bp and CDS: 2127 bp) and had a sequence similarity of 95 and 92% with Ms4's CDS and protein respectively (Additional File 1: Figure S3 and S4). Except MS4_homolog, the next closest protein was only 34.2% similar to MS4 protein which clearly shows MS4_homolog as the only homolog of MS4 in soybean (Data not shown). Further, both the genes were also present as duplicated gene pair in the same segmentally duplicated chromosomal block 245 (chr02-chr14) obtained from Plant Genome Duplication Database (Additional File 1: Table S3) which further confirms the similarity observed in phylogenetic analysis.

Ms4 is highly expressed during early flower development

Since the PHD finger proteins are known to be involved in various functions in plants [20], we looked at the expression of *Ms4* and *Ms4_homolog* across various stages of plant development (Fig. 5). For this purpose, we used publicly available soybean RNA-seq datasets [21, 22]. The comparative expression analysis normalized to root tissue showed significantly higher expression of *Ms4* in early flowering (flower bud differentiation) stages followed by shoot meristematic tissue, 3 weeks old seed, stem and least expressed in root (Fig. 5). Similarly, comparative expression of *Ms4_homolog* normalized to seed_



3w (lowest expression) also showed highest expression in early flowering (flower bud differentiation) stage however, significantly lower than that of Ms4 (Fig. 5). Among the other tissues, $Ms4_homolog$ was also expressed in roots, flower_2 with lowest in 3 weeks old seeds. Among the other 22 genes present in the mapped region, 21 were observed to express moderately across different stages whereas Glyma.02G242500 was constitutively expressed across all the stages analyzed (Additional File 1: Figure S5).

Functional characterization of *Ms4* using Arabidopsis model system

Soybean MS4 (*Glyma.02G243200*) and Arabidopsis MMD1 proteins shared 64% amino acid sequence similarity along with similar conserved domain structure. The Arabidopsis *MMD1* is an essential gene with role in male meiosis and homozygous *mmd1/mmd1* plants are male sterile [16]. Hence, we used the heterozygous Arabidopsis *mmd1* mutant for the complementation with the soybean genes for the functional characterization.

AtMMD1 native promoter driven genomic (Pro_{MMD1} :: gMs4/ mmd1 and Pro_{MMD1} ::gMs4_homolog/ mmd1) and CDS (Pro_{MMD1} ::cMs4/ mmd1 and Pro_{MMD1} ::cMs4_homolog/ mmd1) constructs were used to generate the transgenic plants (Additional File 1: Figure S6). Hygromycin selected T₁ transgenic plants were also genotyped for the homozygous/ heterozygous mmd1 mutant background. Three independent T₁ transgenic lines for each construct were advanced to T₂ generation to select the homozygous mmd1/mmd1 plants and studied for complemented phenotype and microscopic analyses.

The homozygous mmd1/mmd1 plants carrying the Ms4 constructs (Pro_{MMD1} ::gMs4 and Pro_{MMD1} ::cMs4) showed successful complementation of male-sterile phenotype of mmd1 mutant, whereas the homozygous mmd1/mmd1 plants carrying $Ms4_homolog$ constructs (Pro_{MMD1} :: $gMs4_homolog$ and Pro_{MMD1} :: $cMs4_homolog$) did not complement the male-sterile phenotype (Fig. 6; Data not shown for the cDNA complementation). Further, the Ms4 complemented plants produced siliques



with viable seeds (Fig. $6 \,$ e, g, i and k) whereas the siliques of Ms4_homolog complemented plants were similar to the mmd1 mutant (Fig. 6 f, h, j and l). Siliques of the Ms4 complemented plants were observed slightly shorter in size as compared to the Ler-0 plants (Fig. 6 i and k). Since the mmd1/mmd1 mutant produced shorter filaments and pollen-less stamens placing the anthers below the stigma, we further analyzed the flowers from the Ms4 and Ms4_ homolog complemented lines for the differences in the stamens. The microscopic images showed that the flowers of the Ms4 complemented lines produced normal stamens with normal filaments and viable pollens similar to control (Ler-0) whereas the flowers of the Ms4_homolog complemented lines were shorter and pollen-less stamens similar to the homozygous mmd1 mutants (Fig. 6 a-d and Fig. 7 ad). To further confirm the complementation of homozygous *mmd1* defective microspore development, we performed the comparative microscopic analysis of the anther developmental stages of the complemented lines with the Arabidopsis Ler-0 and homozygous mmd1 mutant. The Ms4 complemented mmd1 plants showed microspores with successful tetrad formation which were released successfully from anthers whereas the Ms4_homolog complemented *mmd1* plants showed microspores with shrunken cytoplasm at stage 4 of the anther development, which degenerated and failed to release at stage 8 as observed in anthers from mmd1 homozygous mutant (Fig. 7 e-l). However, there were fewer tetrads and fewer pollens observed in the *Ms4* complemented plants compared to the wild type plants (Fig. 7 e-l).

Discussion

Hybrid seed production and utilization has revolutionized the crop production by exploiting the genetic phenomenon of heterosis/hybrid vigor. This technological advancement directly contributed to an increase in the yield as seen in rice where hybrid rice yield increased $\sim 10-20\%$ over the conventional lines [23]. There is a high potential to exploit the heterosis in soybean; hence, understanding molecular basis of male sterility in soybean mutant lines is critical for the development of high yielding hybrid varieties. Further, insects have shown to be involved in transferring pollen from male-fertile plants to male-sterile soybean plants [24] which avoids laborious manual pollination. Identification and characterization of soybean male-sterile lines are critical for producing hybrid seed at commercial level as well as to understand the mechanism responsible for male sterility. In addition, the genetic male sterility has an advantage over the cytoplasmic male sterility (CMS) that it does not require a dedicated maintainer line to propagate the malesterile line. The major constraint in the utilization of the genetic male sterility for production of pure hybrid seeds is the identification and removal of the male-fertile plants before flowering. Successful pre-flowering recognition of



homozygous male sterile plants requires specific morphological or molecular markers. Male-sterile lines often do not display visible phenotype before flowering hence the gene-specific molecular markers are more practical in the identification and removal of male-fertile plants. In addition, these markers can be used for male-sterile line selection in backcrossing and recurrent selection breeding programs [3]. Further, identification of photoperiod/ thermo-sensitive genic male sterile (PTGMS) lines may provide flexibility in regulating fertility by managing environmental conditions [25]. Hence, developing independent male-sterile lines and understanding the molecular mechanisms will help to deploying a suitable approach for hybrid seed production. In soybean, eleven independent male-sterile lines (ms1, ms2, ms3, ms4, ms5, ms6, ms7, ms8, ms9, msMOS, and msp) generated by various methods (spontaneous, fast neutron irradiation and transposable elements)

have been identified [6–8]. In addition, the availability of the male-sterile lines from diverse genetic backgrounds further facilitates the opportunity for the exploitation of genetic diversity from different lines. Despite the identification of several male-sterile lines in soybean, not much is known about the identity of the genes or molecular mechanism behind the male sterility.

The overarching goal of the present study was to identify and characterize the gene responsible for male-sterile phenotype of the *ms4* mutant. The *ms4* was previously mapped to a 694 kb region on chromosome 2 which harbored 88 genes [6]. The soybean *msp* gene was also mapped in the same region; hence fine mapping to identify the actual causal gene was essential to develop gene-specific molecular markers [6]. It is believed that *ms4* is different from *msp*, as the mechanisms involved in governing male sterility are different [6] indicating that mutations in



independent loci are responsible for the male sterility in these lines. The *msp* is a temperature sensitive male sterile mutant, which displays higher fertility in the hot environment compared to cooler temperatures [14], while *ms4* is not a temperature sensitive mutant. In the present study, we have fine mapped the ms4 locus to \sim 216 kb region that contained 23 protein-coding genes (Fig. 2; Additional File 1: Table S1) and candidate gene was identified based on the functional relevance. The candidate gene (Glyma.02G243200) displayed high homology to Arabidopsis MMD1 which is involved in the male fertility in Arabidopsis [16-18]. The PHD domain protein AtMMD1 has been shown to be essential for proper chromosome condensation during male meiosis [16-18]. The mmd1 mutant fails to produce viable pollen due to pollen degeneration after the tetrad stage, a phenomenon that is also observed in the ms4 mutants, endorsing Glyma.02G243200 as the candidate gene. Sequence analysis of Glyma.02G243200 from ms4/ms4 plants confirmed the presence of a spontaneous mutation resulting into a truncated protein lacking PHD domain (Fig. 3B; Additional File 1: Figure S1 and S2) [17, 18]. The PHD domain has been shown to be critical for MMD1 function in Arabidopsis [16, 18] which is missing in *ms4/ms4* plants due to spontaneous mutation resulting in truncated protein.

Interestingly, phylogenetic analysis of the MS4 protein revealed the presence of another PHD domain protein, MS4_homolog, encoded by Glyma.14G212300 gene on chromosome 14. These two homologs appear to be result of segmental duplication during paleoploidization event (Additional File 1: Table S3). Even though cytogenetically soybean behaves as diploid, there are reports of at least two rounds of whole genome duplication [26, 27]. It is interesting that despite the presence of a homolog, the Ms4 is the only gene that is governing the malesterile phenotype. These results are supported by the expression data which shows that soybean Ms4 expression is significantly higher than Ms4_homolog in most of the tissues, except root (Fig. 5). There are several reports of similar expressional shift (spatial and quantitative) among the remained duplicated genes (paralogs) in plants including soybean [28]. Non-functionality of MS4_homolog can also be explained based on the study in which loss or silencing of approximately 25% of the duplicated genes in soybean were reported since the last duplication event [26, 28]. The Arabidopsis homozygous mutant lines (mmd1/mmd1) complemented with soybean Ms4 (AtMMD1pro::gMs4 and AtMMD1pro::cMs4) showed successful functional complementation by producing viable pollen and siliques with seeds, whereas the Ms4_homolog was not able to complement (Fig. 6).

Since, both Ms4 and Ms4_homolog genes were driven by native MMD1 promoter, the functional characterization clearly showed lack of function for Ms4 homolog, which explains the reason behind the male sterile phenotype of ms4 despite the presence of another homolog. The failure of Ms4_homolog to rescue the male-fertile phenotype could be attributed to the 7% differences in amino acid composition between these two proteins (Additional File 1: Figure. S4). Further studies are needed to understand the reasons for the functional differences. In addition to its inability to produce viable pollens, homozygous mmd1 mutants produce shorter filaments placing the anthers below the stigma (Fig. 6). The complemented Ms4 lines produced normal filaments and viable pollens suggesting that the PHD finger proteins are involved in the filament elongation in addition to the chromosomal condensation (Fig. 6). Overall, our data successfully demonstrated that the spontaneous mutation in Glyma.02G243200 that resulted in premature stop codon is responsible for the male-sterile phenotype of the soybean ms4 line.

Conclusions

Stable male sterility is an important component of an effective hybrid system in self-pollinated crops. This is the first report on map based cloning and characterization of a soybean gene responsible for male fertility. Successful utilization of the male sterility phenomenon requires a complete understanding of associated mechanisms in soybean, which warrants cloning, and characterization of the existing male-sterile lines apart from generating new male-sterile lines. The current study revealed that a similar molecular mechanism is involved in determining the male-sterile phenotype in soybean and Arabidopsis. Information from other plant species can also be utilized to make rapid progress in understanding the male sterility mechanism in soybean through translational research. Employing the model system will result in rapid progress in understanding the molecular mechanism involved in male-sterile phenotypes. A comprehensive understanding of the molecular mechanisms and identification of genes responsible for the male-sterile phenotypes will help to utilize the best system for the production of efficient and pure hybrid seeds in soybean.

Methods

Development of mapping population and molecular mapping

The mapping population was generated by crossing T274H (ms4/ms4) as a female parent with Manchu (Ms4/Ms4) as a male parent. The parental lines were obtained from the USDA Soybean Germplasm Collection [29]. The resulting F₁ plants were self-pollinated to generate F₂ mapping population, which was used for molecular mapping experiment. Fertile/sterile nature of the segregating

population was tested by staining the pollen with potassium iodine solution [30] and also by observing the seed set at the harvest. Upon staining, the fertile pollen turns into dark red-brown while the sterile pollen appears light in color and translucent. The *ms4/ms4* plants did not produce the pods; however, *Ms4/ms4* and *Ms4/Ms4* plants produced the pods with seeds (Fig. 1).

Genomic DNA was isolated from the F₂ plants using CTAB method as described earlier [31]. The sequence information of SSR (Simple Sequence Repeat) markers was obtained from Soybase database [32, 33]. For microsatellite analysis using PCR, 30 ng genomic DNA was used as template in a 10 µl PCR reaction containing 1x reaction buffer (10 mM Tris-HCl, 50 mM KCl, pH 8.3), 2.0 mM MgCl₂; 0.25 µM of each primer; 200 µM of each dNTP and 0.25 units of Biolase DNA polymerase (Bioline, USA Inc., Taunton, MA). The PCR conditions used were as follows: 2 min at 94 °C; 35 cycles of 30 s at 94 °C, 30 s at 58 °C and 1 min at 72 °C, followed by 8 min at 72 °C. The amplified PCR products were separated on a 4% agarose gel. The Mapmaker 2.0 program was used to determine genetic linkages and genetic distances [34]. Marker order was determined at a LOD threshold of 3.0. Linkage calculations were done using the Kosambi mapping function [35].

Amplification and characterization of the candidate ms4 gene (Glyma.02G243200)

Nine combinations of forward and reverse primers were used to amplify the gene from the male-fertile (Ms4/Ms4) and the male-sterile (ms4/ms4) plants (Additional File 1: Table S2A). Long-range PCR was performed using touchdown PCR method with an annealing temperature ranging from 68 °C to 60 °C and extension temperature of 72 °C. PCR products were separated on a 0.8% agarose gel, fragments were excised and purified using gel purification kit (IBI Scientific, Peosta, IA, USA). Twelve pairs of internal primers were used to sequence the full-length Ms4 gene (Additional File 1: Table S2B). The amplified PCR products were sequenced using Sanger sequencing at Functional Biosciences (Madison, WI) and assembled using Vector NTI program (Life Technologies Corporation, Grand Island, NY, USA). Ms4 genomic DNA and cDNA sequences derived from wild type soybean were compared to each other to identify the exon-intron boundaries using Exon-Intron Graphic Maker [36]. Assembled Ms4 genomic DNA sequences from fertile and sterile soybean plants were compared to determine the causal mutation (insertion). Further, the full length and the truncated MS4 protein sequences from the fertile and the sterile soybean lines were aligned using ClustalW, respectively and conserved domains were identified using NCBI Conserved Domains Database (CDD) [37]. Additionally, the physical properties, like molecular weight and pI value of the MS4

protein were also analyzed using ExPASy Compute pI/ Mw tool [38].

Phylogenetic analysis

The amino acid sequence of the MS4 protein was used to identify the homologs in soybean as well as other plant species using BlastP search against the public domain database [39]. The identified 33 MS4 related protein sequences along with the soybean MS4 sequence were aligned using MEGA inbuilt ClustalW alignment tool with standard parameters. Further, the aligned sequences were used to build the maximum likelihood phylogenetic tree using MEGA v6.06 software [40] with 1000 times boot strap replications. Other parameters used for tree construction were: Substitution type: Amino acid; Model: Equal input model; Rates among sites: Uniform rates; Gaps/missing data treatment: Partial deletion; Site coverage cutoff: 95%.

Developmental stage specific expression analysis

To perform the soybean *Ms4* and *Ms4_h* expression analysis, 17 publicly available RNA-seq datasets related to various vegetative (cotyledon (2), leaf (3), stem (2), root and shoot_meristem) and reproductive (flower (5), pod, seed and pod_seed) stages of soybean development were downloaded from the NCBI SRA database (accession PRJNA238493) (Additional file 1: Table S4). All the datasets were analyzed using RNA seq analysis tool of CLC Genomics workbench. Expression data related to *Ms4* and *Ms4_homolog* was extracted, and fold change was calculated using the tissue with minimum expression as control. A heat map was plotted for the expression data of 22 other genes present in the mapped region using the online tool Morpheus (https://software.broadinstitute.org/morpheus/).

Complementation vector construction

Soybean plants (Dyna-Gro S56RY84) were grown under controlled greenhouse conditions at 28 °C temperature. Leaves and flower buds were collected from the flowering stage plants and flash frozen using liquid nitrogen and stored at - 80 °C. The genomic DNA from leaf tissue was extracted using DNeasy kit (Qiagen, Valencia, CA USA). RNA from flower buds was extracted using the plant spectrum total RNA isolation kit (Sigma Aldrich) and simultaneously treated with on column DNaseI (Sigma Aldrich) using standard protocol. Further, 1 µg of DNaseI treated RNA was used for the cDNA synthesis using the SuperScript II reverse transcriptase (Invitrogen). The Ms4 (Glyma.02G243200) and its homologous gene (Ms4_homolog; Glyma.14G212300) sequences obtained from phytozome soybean genome sequence data (https://phytozome. jgi.doe.gov/pz/portal.html) were used to design primers to amplify the full-length sequence for cloning and complementation studies (Additional File 1: Table S2C). Since both the genes share very high sequence similarity at the 5' and 3' ends, same primer pair was used to amplify both the genes. To drive the expression of both the genes, Arabidopsis native MMD1 promoter was amplified using the specific primer pair (Additional File 1: Table S2C) and cloned into pMDC32 expression vector using Sbfl and AscI restriction enzymes (New England BioLabs, Ipswich, MA, USA). The Ms4 and Ms4_homolog genes (genomic DNA fragments) were amplified using NEBNext® High-Fidelity 2x PCR Master Mix (New England BioLabs, Ipswich, MA, USA) with the PCR condition as follows: 3 min at 98 °C; 30 cycles of 30 s at 98 °C, 30 s at 58 °C and 2 min at 72 °C, followed by 10 min at 72 °C. The amplified products were separated on a 1% agarose gel and DNA fragments of ~ 3.1 kb (gDNA) and ~ 2.1 kb (cDNA) were gel purified using a GeneJET Gel extraction kit (Thermo Fisher Scientific, Carlsbad, CA, USA). The purified PCR products were cloned into the expression vector pMDC32 containing Arabidopsis MMD1 native promoter using AscI and PacI restriction enzymes and T4 DNA ligase (NEB, Ipswich, MA, USA). The ligation mix was transformed into DH5- α chemical competent Escherichia coli cells. Plasmid was extracted from the colony PCR positive colonies using GeneJET Plasmid Miniprep kit (Thermo Fisher Scientific, Vilnius, Lithuania) and further confirmed by restriction digestion with AscI and PacI enzymes. The Ms4 and Ms4_ homolog specific constructs were confirmed using the restriction digestion with HindIII which is only present in the Ms4 and not in Ms4_homolog. The sequencing confirmed genomic (AtMMD1:gMs4 and AtMMD1:gMs4_ homolog) and CDS (AtMMD1:cMs4 and AtMMD1:cMs4 homolog) constructs were transformed into Agrobacterium tumefaciens using standard transformation protocol and PCR positive colonies were used for plant transformations.

Plant transformation

Complementation studies were performed using the male meiocyte death 1 (mmd1) mutant of Arabidopsis thaliana ecotype Landsberg erecta (Ler-0) [16]. Since, the homozygous mmd1 is male sterile, only heterozygous (MMD1/ mmd1) plants can be transformed. The mutant plants were genotyped for heterozygous nature before transformation using gene-specific forward and dissociator (Ds) specific reverse primer pairs as described previously [16]. The genotyped mmd1 heterozygous plants were transformed with soybean Ms4 and Ms4_homolog genomic (AtMMD1: gMs4 and AtMMD1:gMs4_homolog) and CDS (AtMMD1: cMs4 and AtMMD1:cMs4_homolog) complementation constructs using A. tumefaciens (GV3001) mediated floral dip transformation method [41]. The T_1 transgenic seeds were grown on Murashige and Skoog (MS) media with Hygromycin (50 μ g/ml) to select the transformants. The selected T₁ transgenic plants were advanced to T₂

generation to isolate homozygous lines which were used for further functional characterization.

Microscopy analysis

Mature flower and silique samples were collected from the *Ler-0*, homozygous mutant (*mmd1/mmd1*), *Ms4* (*mmd1/AtMMD1:gMs4* and *mmd1/AtMMD1:cMs4*) and *Ms4_homolog* (*mmd1/AtMMD1:gMs4_homolog* and *mmd1/AtMMD1:cMs4*) homolog (*mmd1/AtMMD1:gMs4_homolog* and *mmd1/AtMMD1:cMs4_homolog*) homozygous transgenic lines to image stamens, pollens using light microscope BX-41 (Olympus, MA, USA) and siliques using SZ61 stereomicroscope (Olympus, MA, USA). Anthers were separated from the flower and stained with I₂KI solution [42]. Dark red color staining, an indicator of live pollen was used to compare the pollen from *Ler-0*, mutant and complemented lines. Microspore development was studied at stage 4 and stage 8 of the anther development, using light microscopy [43].

Additional file

Additional file 1: Figure S1. Sequence comparison of the genomic DNA sequences showed an insertion of a single nucleotide "A" in the 3rd exon of the Ms4 gene (Glyma.02G243200) in the male-sterile (ms4/ms4) line. Green color boxes represent exonic sequences. Insertion mutation is shown by black color box. Figure S2. Amino acid alignment showing a frameshift mutation resulted in an early stop codon in ms4 sterile mutant. The resulting truncated protein lacks the Plant Homeodomain (PHD) which is otherwise present at the C-terminus of native MS4 protein. Figure S3. Sequence comparison of the soybean Ms4 and Ms4_homolog coding sequences. Figure S4. Similarity comparison of the soybean MS4 and MS4_homolog amino acid sequences. Figure S5. Heatmap representing the expression of the remaining 22 genes present in the mapped region. Figure S6. Constructs used for the complementation purposes. (A) Arabidopsis MMD1 native promoter driven genomic and CDS Ms4 constructs. (B) Arabidopsis MMD1 native promoter driven genomic and CDS Ms4 homolog constructs. Table S1. A list of 23 predicted genes in the ms4 region flanked by BARCSOYSSR_02_1515 and BARCSOYSSR_02_ 1528. Table S2. Details of primers used in present study for various purposes. Table S3. Duplication gene list of syntenic block 245 between soybean chr02-chr14 with Ms4 and Ms4_homolog as duplicated gene pair (highlighted in yellow). Table S4. List of SRA files used for the expression analysis of Ms4 and Ms4_h. (PDF 672 kb)

Abbreviations

BLAST: Basic local alignment search tool; CDS: Coding sequence; MMD1: MALE MEIOCYTE DEATH 1; MS: Male sterile; PHD: Plant Homeodomain

Acknowledgments

We thank Dr. Christopher A. Makaroff for generously providing seeds for the $\mathit{mmd1}$ mutant.

Authors' contributions

SWT fine mapped the *Ms4* locus, cloned the *Ms4* gene, worked on transformation experiment, performed the analysis and wrote the first draft of the manuscript. KMR performed the bioinformatics analysis, conducted transformation in Arabidopsis and performed analysis. AR and VKB helped in bioinformatics analysis. RGP generated the mapping populations. DS and VM conceived the idea, helped in interpreting results and supervised the findings of the work. SWT, KMR, DS, AR, VKB and VM contributed towards writing the manuscript. SWT, KMR, DS, AR, VKB and VM have read and approved the manuscript (RGP deceased).

Funding

Not applicable.

Availability of data and materials

We have used publicly available RNA-seq data from the NCBI SRA database BioProject: PRJNA238493. More details are available in Additional file 1: Table S4.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Fiber and Biopolymer Research Institute, Department of Plant and Soil Science, Texas Tech University, Lubbock, TX 79409, USA. ²US Salinity Laboratory (USDA-ARS), Riverside, CA 92507, USA. ³Wisconsin Institute of Sustainable Technology, University of Wisconsin-Stevens Point, Stevens Point, WI 54481, USA. ⁴Department of Agronomy, Iowa State University, Ames, IA 50011, USA.

Received: 20 April 2019 Accepted: 15 August 2019 Published online: 28 August 2019

References

- Hartman GL, West ED, Herman TK. Crops that feed the world 2. Soybean—worldwide production, use, and constraints caused by pathogens and pests. Food Security. 2011;3(1):5–17.
- Lewers KS, St. Martin SK, Hedges BR, Widrlechner MP, Palmer RG. Hybrid soybean seed production: comparison of three methods. Crop Sci. 1996;36:1560–7.
- Lewers KS, Palmer RG. Recurrent selection in soybean. In: In: Plant Breed Rev. Inc: Wiley; 2010. p. 275–313.
- Ott A, Yang Y, Bhattacharyya M, Horner H, Palmer R, Sandhu D. Molecular mapping of D1, D2 and ms5 revealed linkage between the cotyledon color locus d2 and the male-sterile locus ms5 in soybean. Plants. 2013;2(3):441.
- Speth B, Rogers JP, Boonyoo N, VanMeter AJ, Baumbach J, Ott A, Moore J, Cina T, Palmer R, Sandhu D. Molecular mapping of five soybean genes involved in male-sterility, female-sterility. Genome. 2015;58(4):143–9.
- Yang Y, Speth BD, Boonyoo N, Baumert E, Atkinson TR, Palmer RG, Sandhu D. Molecular mapping of three male-sterile, female-fertile mutants and generation of a comprehensive map of all known male sterility genes in soybean. Genome. 2014;57(3):155–60.
- Cervantes-Martinez I, Xu M, Zhang L, Huang Z, Kato KK, Horner HT, Palmer RG. Molecular mapping of male-sterility loci *ms2* and *ms9* in soybean. Crop Sci. 2007;47(1):374–9.
- Cervantes-Martinez I, Sandhu D, Xu M, Ortiz-Pérez E, Kato KK, Horner HT, Palmer RG. The male sterility locus *ms3* is present in a fertility controlling gene cluster in soybean. J Hered. 2009;100(5):565–70.
- Albertsen MC, Palmer RG. A comparative light- and electron- microscopic study of microsporogenesis in male sterile (*MS*₁) and male fertile soybeans (*Glycine max* (L.) Merr.). Am J Bot. 1979;66(3):253–65.
- Graybosch RA, Palmer RG. Male sterility in soybean (*Glycine max*). II. Phenotypic expression of the *ms4* mutant. Am J Bot. 1985;72(11):1751–64.
- Jin W, Horner HT, Palmer RG. Genetics and cytology of a new genic malesterile soybean [*Glycine max* (L) Merr.]. Sex Plant Reprod. 1997;10(1):13–21.
- 12. Skorupska H, Palmer R. Genetics and cytology of the *ms6* male-sterile soybean. J Hered. 1989;80(4):304–10.
- Palmer R. Genetics of four male-sterile, female-fertile soybean mutants. Crop Sci. 2000;40(1):78–83.
- Stelly DM, Palmer RG. A partially male-sterile mutant line of soybeans, *Glycine max* (L) Merr.: characterization of the *msp* phenotype variation. Euphytica. 1980;29(3):539–46.
- Delannay X, Palmer RG. Genetics and cytology of the *ms4* male-sterile soybean. J Hered. 1982;73(3):219–23.
- Yang X, Makaroff CA, Ma H. The Arabidopsis *Male Meiocyte Death1* gene encodes a PHD-finger protein that is required for male meiosis. Plant Cell. 2003;15(6):1281–95.

- Andreuzza S, Nishal B, Singh A, Siddiqi I. The chromatin protein DUET/ MMD1 controls expression of the meiotic gene *TDM1* during male meiosis in Arabidopsis. PLoS Genet. 2015;11(9):e1005396.
- Wang J, Niu B, Huang J, Wang H, Yang X, Dong A, Makaroff C, Ma H, Wang Y. The PHD finger protein MMD1/DUET ensures the progression of male meiotic chromosome condensation and directly regulates the expression of the condensin gene *CAP-D3*. Plant Cell. 2016;28(8):1894–909.
- Gasteiger E, Hoogland C, Gattiker A, Se D, Wilkins MR, Appel RD, Bairoch A. Protein identification and analysis tools on the ExPASy server. In: Walker JM, Totowa NJ, editors. In: *The Proteomics Protocols Handbook*: Humana Press; 2005. p. 571–607.
- 20. Sanchez R, Zhou M-M. The PHD finger: A versatile epigenome reader. Trends Biochem Sci. 2011;36(7):364–72.
- Shen Y, Zhou Z, Wang Z, Li W, Fang C, Wu M, Ma Y, Liu T, Kong L-A, Peng D-L, et al. Global dissection of alternative splicing in paleopolyploid soybean. Plant Cell. 2014;26(3):996–1008.
- Golicz AA, Singh MB, Bhalla PL. The long intergenic noncoding RNA (LincRNA) landscape of the soybean genome. Plant Physiol. 2018;176(3):2133–47.
- Zhou H, Zhou M, Yang Y, Li J, Zhu L, Jiang D, Dong J, Liu Q, Gu L, Zhou L, et al. RNase ZS1 processes *UbL40* mRNAs and controls thermosensitive genic male sterility in rice. Nat Commun. 2014;5:4884.
- Palmer RG, Gai J, Sun H, Burton JW. Production and evaluation of hybrid soybean. In: In: *Plant Breed Rev.* Inc: Wiley; 2001. p. 263–307.
- Frasch RM, Weigand C, Perez PT, Palmer RG, Sandhu D. Molecular mapping of 2 environmentally sensitive male-sterile mutants in soybean. J Hered. 2011;102(1):11–6.
- 26. Shoemaker RC, Schlueter J, Doyle JJ. Paleopolyploidy and gene duplication in soybean and other legumes. Curr Opin Plant Biol. 2006;9(2):104–9.
- Schlueter JA, Lin JY, Schlueter SD, Vasylenko-Sanders IF, Deshpande S, Yi J, O'Bleness M, Roe BA, Nelson RT, Scheffler BE, et al. Gene duplication and paleopolyploidy in soybean and the implications for whole genome sequencing. BMC Genomics. 2007;8:330.
- Lynch M, Conery JS. The evolutionary fate and consequences of duplicate genes. Science. 2000;290(5494):1151–5.
- 29. Germplasm Resource Information Network National Plant Germplasm System [https://www.ars-grin.gov/npgs/]. Accessed 19 Aug 2019.
- Jensen WA. Botanical histochemistry: principles and practice. San Francisco: W. H. Freeman; 1962.
- Sandhu D, Gao H, Cianzio S, Bhattacharyya MK. Deletion of a disease resistance nucleotide-binding-site leucine-rich- repeat-like sequence is associated with the loss of the Phytophthora resistance gene *Rps4* in soybean. Genetics. 2004;168(4):2157–67.
- Song QJ, Marek LF, Shoemaker RC, Lark KG, Concibido VC, Delannay X, Specht JE, Cregan PB. A new integrated genetic linkage map of the soybean. Theor Appl Genet. 2004;109(1):122–8.
- SoyBase: Integrating Genetics and Genomics to Advance Soybean Research [https://www.soybase.org/dlpages/index.php]. Accessed 19 Aug 2019.
- Lander ES, Green P, Abrahamson J, Barlow A, Daly MJ, Lincoln SE, Newburg L. MAPMAKER: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. Genomics. 1987;1(2):174–81.
- Kosambi DD. The estimation of map distances from recombination values. Ann Eugenics. 1943;12(1):172–5.
- WormWeb: Exon-Intron Graphic Maker [http://wormweb.org/exonintron]. Accessed 19 Aug 2019.
- NCBI Conserved Domains Database [https://www.ncbi.nlm.nih.gov/ Structure/cdd/wrpsb.cqi]. Accessed 19 Aug 2019.
- ExPASy: Bioinformatics Resource Portal [http://web.expasy.org/compute_pi/]. Accessed 19 Aug 2019.
- NCBI Basic Local Alignment Search Tool [http://blast.ncbi.nlm.nih.gov/Blast.cgi]. Accessed 19 Aug 2019.
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. Mol Biol Evol. 2011;28(10):2731–9.
- 41. Clough SJ, Bent AF. Floral dip: a simplified method for agrobacteriummediated transformation of *Arabidopsis thaliana*. Plant J. 1998;16(6):735–43.
- 42. Sulusoglu M, Cavusoglu A. *In vitro* pollen viability and pollen germination in cherry laurel (*Prunus laurocerasus* L.). Sci World J. 2014;2014:7.
- Horner HT, Palmer RG. Mechanisms of genic male sterility. Crop Sci. 1995; 35(6):1527–35.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- · thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

